



g. x sin

one thought to
be due to this.
- Quincke's edema
Etiology

As they are
to some extent
disturbance such
as getting fish,
crops, pop. exposure
for a most any
thing. Anticoin is
called metacere
+ gives, then it
me. due to ingestion
from edema Quinine
crops, dose, the
come to prep-
in fact most any
of drugs may give
~~first to the~~

~~It may be~~

use of
injection of diphtheria
antitoxin of any
granulation may
give rise to it. &
if using of a needle
or of an instrument may
cause it. the
wearing of shoes
under clothing
may give rise to it
in those predisposed
It is very difficult
to locate & cause so
our chronic varieties
of me, orsted,
with inflammation
gout, & ist poritis
with Bright's dis.

The larvae that

merely shut id
those lesions of
great importance
they may reach
size of other eggs.
So the egg lesions
stretch & if on
side of face.
On inst. & tongue
suddenly wells
the whole ridge,
that me. the
mouth won't
hold it. These
egg lesions usu-
ally become give
rise to swelling
& discomfort
^{thought to be}
some spider bite.

In babies that
suffer from constipa-
tion or other gastro-
intestinal derangement
there is fully a
small pro-
portion who take
nothing at all &
are very resistant
to treatment.

meat!

I very simply
you forget at
depending on time
before it leaves
Now on electric
time physical
attack & shot.
other wire a bush

without the size

about the size of
beaver or spirit
head not more,
3 or 4 in across
and is usually
on a neck about
the size of your ^{little} finger
mid. of length.
is symmetrical.
V points of hds. &
roes of feet are
not involved.
V marks of fine
attached. Where V
is in the plowing
became involved
right of impression
the very distressing
V person not involved

1/2 tsp 1 mark

solution of pyrazine
administered
about 90%
this will relieve
it. A more
intolerable pyz,
require some
antipruritic,
as the sol. of too
much of carbonic
acid or ether or
comp. Carbonic
acid is best antipruritic we have
It is used in
1-2% depending
upon strength
of agent.
An alkaline sol.

we stay looking on

Latent comes
out very suddenly
the itching is usually
very intense & c
itching milder
in a few hours to a
few days. In ^{some} ~~some~~
instances it may
last for several
weeks. due to a de-
pendent of success-
ful crops of eruption
& eruption is often
very wide spread.
At times it is
confined to only
a few wheals,
& individual
lesion is small.

the dog dived on
off on a hot bath
with should feel
of cooling soda
will really
give great deal
of relief - the coal-
tion which is anti-
pyretic taken in
about 5 gr. doses
will. They give
great relief.
There is 1 woman
one present they
should be given
id of and this will
prevent v & con-
dition. If we had
to discover v cause

h

the fine structure
of the tissue

Inflammation Thrombosis

Is a stage of
the blood
by presence of
fibrin, that of a
very pronounced
state quite
intensely. & when
in condition is
usually a flat
topped slight elevation
& borders of rock are
hyperemic white &
usually pale.
It due to an edema
of very slight ones
it bed to find for
& O. & when in

~~Constitution~~
in spite of every
thing you do the
cough will per-
sist. Chronic

cough, and some
most distressing
things to me. I
see. If the
cough is present
give appropriate
remedies. If you
are bright, do
relieve as far
as possible.

In a majority of
cases a brisk
purgative & an applica-
tion of ointment to relieve
itching should
be used.

Frederick, Nov. 24 - 08

On by means of
some south by
lotion kept con-
stantly applied,
where condition
is somewhat there
is considerable
impl. & weeping
in addition to
that above you
should keep
your reported
by means of some
soft part,
Engthman Intention
So get well away
kidding when treated
in this way.

Nov 20

Lecture, Nov. 24-08

Erythraea multiformis
Eryth. mult. is
char. by imper-
fect condition associated
with papules &
tubercles, vesicles
& bullae. *E. ery-*
thraea is symmetri-
cally distributed
there are a no. of
varieties of *mult.*
Eryth. mult.

depending upon
color of the skin
present. & contours
Most fr. varieties
it is popular & tuber-
culous.

A no. of dif. varieties

Neck in
creased by
upward skin

neck in
creased by
upward skin
The neglected case
Dyflon in the
Oiled some degree
prod. As smoke
there is reddening
a burning and
then or perhaps slight
itching. It is
counted as very
marked.
Treatment

Consists
in cleanliness &
use of some
rooting dressing
found much of
it & stolen

Lesions are present
in some cases but
as a rule one
variety predominates
the lesions of E.M.
come out ^{very} typical & develop
on a rule mature
very slowly. Lesions
are for first bright
red & later course
of a dozen two days
become of a scarlet
color. Papules
vary in size
from a ^{small} ~~open~~ head to
a split pea. The
tubercles vary
from a split pea
on up to almost
langer.

Alfred Wilson

I opposed it.
I think more
in it was
nothing so bare
persons I say
and in getting
elect. I say
I mean with
pendulous breasts
occurring between
I breast & chest
was in those
who are pretty
fleshy & have been
so covered by
fat about the
you will find
it in the
muscles. I say
in lobes about

the vesicle

When v tubercles &
popules are present
= E.T. or E.P.
If tubercles are present = Ength. tubercula
If popules are present = Ength. populeosa

So the influ.
change is greater
than to prod tuberc.

or popules. The
youngst a vesicle
or bullae than
we speak of it as
Bullae or vesicular
Ength.

It. E. M. begins
or abroad plot
popule about v
size of a dime. This
lesion may clear
up in center &
spread at periphery

During Idiopathic
rheumatic truma-
tic erythema.

✓ Erythema due
to sun burn or to
frost bite, the
symptomatic erythema
that occurs with
would be blushing
also one due to
some gastro intestinal
trouble or to
irritation of some
kind.

Erythema Intestin-
go. Only spoken
of as chapping.
Is a hyperemic condition
not produced by rubbing
together of

~~Over 2 mm~~

giving rise to annular
Enyth - Enyth - annular
there are, reversal
of these annular
lesions present
close together &
these lesions may
coalesce the lines
of coalescence disappear
or subside leaving
a lesion of an
inner - original
outlet = Enyth.
marginal or
Enyth ^(cystic) pyrotonin.
At times Enyth. M.
begins as one of these
sub. annular lesions
& concentric with
that sub. annular

Lesson on the

Hyperemias.
The hyperemias are
those conditions in
which there is some
hyperemia of the skin
not associated with
any infiltration or
exudation. I divide
my line between
✓ hyperemia and
✓ inflammation not
very well drawn.

Erythema simplex
is a idiopathic
or symptomatic.
✓ Idiopathic is due
to some extrinsic
✓ Symptomatic and
due to something
that acts internally.

~~commonest type~~

lesion another
annular lesion
develops & perhaps
another one. (3) (4)
After a older lesion
begins to fade a little
by & time next
concentric lesion
appears & next
lesion & same
so that you have
reverse concen-
tric lesion of
somewhat dif-
ferent color. become
of a diff color we
speak of this as Eryth-
Tris.
At times Eryth will
begin as a black & then

P.C. can also be
destroyed by some
of a fine comb.
This is a contagious
condition. It can
transmit itself by close
contact with those
affected or by
means of the hands
of those infected.
Very common in school
children especially
in those commu-
nities where clean-
liness is neglected
more or less.
Spec

P. G. can see be

with some centers
a ring of vesicles
will bech and
that black & another
ring of vesicles will
often around
so that you may
get from 2 to
concentric rings
of these & black
they differ some
what in color due
to age. As spoken
of as Herpes Iris
or Eryth-Herpes Iris.
they are very
very rare forms
of E. M. & popular
+ tubercular varieties
and most an.

extinguish the
flames can be
removed by p.
working with 10
H₂O or with
dense acetic acid.

Dye is very much
shaded & is a great
deal of rusting. It is
better to cut & then
not removing
crusts. Or if
you want cut
these crusts.

worked out completely
with sweet oil &
fender penum &
like & like at
some time.

~~up to~~

V lesions of eye,
develop suddenly
& mature as a
rule in from a
few to 24 hrs.
& individual lesions
as small as seeds,
in from 3 or 4 days
to week. at first
bright red & V
color deepens to
oblivious to scarlet
this bluish tinge &
rather characteristic.

In course of 7-10
days lesions disappear
leaving behind a
stain to mark its
site. ~~for a while~~
& ~~for a while~~

From 20 to 25

27 Infe. case. in
Mollusca.
testis

Very disappointing
try. Can be performed
Abundant a dense
with = ^{olive} ~~oil~~
oil applied morn-
ing & night for
3 or 4 days will de-
stroy lice. Old
1st. Staphylinia.
1st. 3rd
occure but die.

will also destroy
lice. 1/2 oz. shot
1/2 oz. applied
morning & night
for reverse dogs will

Location at corner
E. W. Dr &
development &
back of hands
& ^{up} to
elbow - on
& down of feet
to knees.

Not. imply - va-
riation at back &
plumage & color
of. then in very
fine. sticks &
neck & face &
to the time.

parent one think
It shows a decided
preference for
horse. of of hands
and arms up to elbow and

I believe the best way
to keep and improve the
mind

give rise to life.
Of neighboring
sympathetic glands.
Especially the glands
of the about the thyroid
process and others
they secrete.

On the very worked
cores the skin will
become matted together
in part mulling
events beneath
back. There are some
thick pus almost
working with the
there the nose &
neglected. In the
majority of cases
the matted gives
rise to no life.

Give rise to
v down for feet up to
the legs and up to the
thoracic part
of any of v body
+ may give attack
v m. In some
instances it may
be confined to trunk
v subjective symp.
of E.M. are often
dreadful - there may
be slight itching or
burning. There
are some few
constitutional syms. that
precede the onset. a
last for a time after
v outbreak. m. has
a slight malaise
+ relative temp.
rise.

General Summary

they attach their
egg to a spot of
hair near the
the base of the
3-10 days. Head
lice give rise to
marked itching
In some instances
it may give rise
to very marked
secondary eczema.
In fact, any pustular
eczema of the body
should be looked
to perfect presence
of the especially
when more exten-
sive on the occipital
region. These
cases of infestation

May 2nd 1892

slight joint
symp. Rheumatic
heat in muscles
& joints these
rheumatic joints
be not infrequent.
Rarely there is
symp. more rather
sharp - & in some
instances E. M.
may occur or one
of frequent
some very good
constitutional disorder.

In a majority of
instances however
a general symp. are
very slight. Most
p. being & pains
in joints & muscles.

Lesson 11

and 1/2 mts.
most any other
parasitoid will
do slightly,

Lecture Nov 17th 08.

Pelionosis capitis -
Common of 1 lb.
Is prod. by 1 presence
of 1 lb. house on
roofs. more com.
in places than
in the children
than adults. Is
said to be more
com. in white
people than in
negro. Is more com.
in the tropical
region than elsewhere

and the young
E. M. usually
goes out to roost
at night, to a whist
there are many
times or out crop-
ping, that will
keep it going for
2 or 3 hours and
in some rare in-
stances it may
keep occurring &
make it appear as
if E. M. occurs a little
frequently.
In most cases
however, it is
~~difficult~~ to find
it occurring
after 4 o'clock.

then they deposite
eggs on which
there very intense
itching they are
prey, gotten from
several intercourse
and from public
closets,
Treat.

Mercurial
is very efficient
and very costly.
Hydroxide 1-5000
will destroy them
* Na hypochlorite
sol. 1/3 to 1/2 will
destroy them it
is well to wash
bath with little
beine acid to get

from the long

h. between ages
of 25 & 30. occurs
most often during
changeable weather
of spring & fall
is frequently associated
with gonitis
& is associated with
rheumatism, gonitis,
Bright's disease & dis-
order of menstruation
may follow adminis-
tration of copaiba
K & L correction.
Is believed to be
most often due to
gonitis with drainage
inward. Was held
or once to be an
organic neurosis.

It has been noted

I + to find some
systemic poison
disturbance & control
of various other
bacteria, & blood, sugar
of skin. It is
evident of deeper
foci of skin
involved with some
some round cell
infiltration & some
exudation of cells.
Treat.

Everyone
will get well if
you leave it alone
or once a Mercurochrome
or white purg should
be given. Stop &
rheumatic position

~~the~~ ~~confront~~

✓ Course of these
state colored potatoes
it not exactly
known but is sup-
posed to be caused
by something injec-
-ed into ✓ body by
while I you will
bake ✓ clothes you
may come in
this way there
they have been
burnt for a long
time you may
see sulphur pits
to kill those on
✓ fine hairs.
Where you see these
long scratch marks

1 Come / Stone
trouble - you
should go at once
upriver to Bodin
noting the. Of
which I send
you to give some
information especially
by collecting and
a little material.
You find a certain
amount of water
alkaline both
from the dist.
get better in
7-10 days.

~~Ding~~

f. Chonnet.
location rapid
open. then
high red color

~~Long scratches~~
~~Long scratches~~
+ neck and you
have long scratch
marks you may
inspect body ^{size} ~~color~~
marked patches
these scratch marks
as you may find
little punctate be.
+ these are where
lice have been
biting. Where the
lice have been on
for sometime you
may have little
dots colored or grey-
ish patches. When
a pet has had them
for long time these
marks may be quite
numerous.

which is your

of later & bluish
color ~~than~~ the
~~lower than~~ the
rapidity of the occur-
rence of the lock
of subjective symptoms

Lecture Nov. Dec. 12th 08.

Eryth. nodosum.

Is an ac. in fland. dis-
chor. by & develop-
ment of older pro-
nities. & dis. m. begins
with malaise or with
considerable fever.

Rheumatic pain are
nearly always present,
but they are quite in-
tense. They may
involve either & usually

25' 2000 3000 4000

When they get
very populous and
there is not room
for all of them in
a room of clothes
they may lay their
eggs on a fine clothes
line. In body.
A body louse sometimes
thinks a head louse
and does not
attack a head,
A body louse causes
intense itching &
is especially about the
neck & back of shoul-
ders. If a patient
comes to you com-
plaining of itching
about the shoulders

John Thompson

or joints or both, & the
skin joints are not
swollen. reminding
one somewhat of re.
gouty rheumatism.
Usually in from a few
hrs. after a foot begins
to feel bad or aching
this. Some spots appear
in plain & swelling
very takes place in
these some places,
giving rise to a node
on a common swelling
in size from a spot
pea to a hen egg.
Swelling grad. fades
into & subsiding this.
~~these~~ ^{these} have no
sharp line of de-

It very seldom
attacks children
& those who are
strong in body.
But it may do
so.

Body lice are very
peculiar to vagabond
clown troupes etc.
There are ^{lice} ~~also~~ usually
five in a clothing
& go down to the
bottom of feet.
They lay their eggs
in the seams of
clothing. They may
be on the seams of
waist and in
waist band of
garments.

March 18

monotonous they
are ⁱⁿ symmetrically
disposed being long
always placed ✓
on the side of the leg;
15 they are scattered
to 2 thighs & some
to the arm. present
on the upper extre-
-mities & so they will
be present on the arm
but this is very rare.
these tumors usually
have these long dis-
posed parallel to the
long axis of the limb
they vary in num-
ber from 4 or 5 to
1 - 3 doz. I begin-
ning they are rather

taken my ^{plithyrosis}
Pediculosis = ^{plithyrosis} ~~plithyrosis~~
thiasis
rosis = Lice

there are three
~~varieties~~ ^{varieties} Note that
affect human
body, i.e.

1. Body louse =
Pediculus Vestimentorum

2. Pediculus Capitis.

3. Pubis =
crotch louse.

Pediculosis Vestimentorum

this condition
attacks & aged and
diseased health
is not quite up to
work.

bright red in color
+ are painful. As time
goes on the color deep-
ens until they be-
come violet or
bluish ~~color~~ just as
a bruise does, and
as they get older
they soften so that
as you can almost
get fluctuation.
They do not
break down save in
very rare instances.
they become absorbed
& their pitte is marked
by a stain which grad-
ually disappears. The individ-
ual runs its course
from 1 week to 10 days.

There are more than
Naphthomelplite
13 to 15 oz. Applied
twice daily will
destroy the parasite
H. c. line, used
where it is not
wide spread your
remedies may prod.
a dermatitis after
✓ scabies are des-
troyed for that
reason & treat. only
lasts for 2 or 3 or 4
days. If after ✓
prod. has treated
for a few days and
it appears that
it is not cured
✓ treat. should be

There are more than
one crop which
prolongs its life for
3 or 4 yrs. It occurs
more early in females
than in males, &
its time of preference
is between ages of
18 & 30. In a group
of 168 cases only 10
cases occurred after
age of 40. The persons
have but little to
do with the time
of occurrence, but
at any difference
of temperature ju.
in spring sometimes
& lesion of Crypt.
nodosum on it.

Overcast with clouds

Nephthol + Balsam

Opium

Beta nephthol

15 gr to 1/23. +

Balsam Opium

13 to 1 oz of base line

you may use
nephthol with
this. If the pot.
object to r. and
the pot. may take
about a teaspoon
of dry washed sugar,
into his plain
and then sprinkle
the same and.

between 1 sheet
and will come
— about 1 wh.

Notch & Bottom

associated with lesions
of Cerebral multipolar
and many there
is some thing in
common between
V cones, of V two
troubled. Many
of observers conclude
that eighth nerve
is closely related to
neurotism in
its course. It has
also been seen to be
affected with various
conditions. including
dementia, various
nervous disorders
& some of nervous
diseases. It is serious
but troubles follow

only minor 2

3 or 4 successive
days. Rub in thoroughly
✓ sulphur may
cause a dermatitis,
this will cure
all ✓ sores of itch.
Use Vaseline to mix
✓ sulphur with.
~~we can~~ ✓ some
under clothes for
3 or 4 days. ✓ pat.
should take a bath
put on clean under
clothes & disinf. ✓
under clothes &
bed clothes.

In some very
delicate skin, sulphur
is too irritating
to use than beta

3 or 4 massive
E. Wilsoni just
when they may in the
time. It is thought
by some E. Wilsoni
is related to some
miss. but is not
very probable,
Pottery

It is that of a
round cell in the
trunk. covered with
some resin and
freq. bld. occurring
in the deep layers of
the trunk and it is the
deep mud and this,
after resolution
the series under-
go & especially the
some changes as

These figures

and then upon
presence of the
the timber or com-
-rains

Treaty

Seems satis-
factory. Sulphur
and is not every
elegant preparation
or most satisfactory
for dis. As to my
ferry from scales
is instructed to
take about both in
road & water so as to
remove all foreign
material but then
to apply on a
contributing 13 Sul-
phur to 100 lb of
for

June 1

+ other things being
equal is more and
for those whose health
is run down for
any cause but,
that does not make
much difference.
Iitch water likes
to attack the thin
skin portion and
where there is much
dilatation you
may often find
to get the
diagnosis

is fairly early
+ depends upon
character & ^{location} ~~area~~
of erupt. whether
prolonged, just or over

Treat.

— I am entirely
satisfied with the
committee. Since the
letter corrected by
appropriate remedy,
the policy votes are
most appropriate
in remedies, we have
it. they are very
correct upon
it. The plan is
good some of them
we have advised
me of as
St. the plan does
seem that it re-
quires opening to
control it, &
local remedies are

✓ Charact. determined
Is visible to ✓ naked
eye as small Meck
Is contagious and
requires rather in-
imate association
for ✓ contagion to be
carried as
~~known~~ as a rule,
such as wearing ✓
some clothing or sleep-
ing between ✓ some
bed clothes,
scabies more easily
attracted ✓ cooler
but is by no means
confined to that
class. Is much worse
in winter time
when people don't
bathe much

1 Chlorostilbe - graysi

usually success
less than 10 days
~~usually~~ but appear
cousin of Boie
said about 2 years
good. I put should
be kept in very
quiet, &
I have diet
regulated. This con-
dition generally
runs its course
in 7-10 days. Recur-
rences may occur
but are rare.

E. nodosa is not
very common it
is very rare.

Pelagra, head up.

~~generous amount~~

~~amount~~ 3

itch mite me.
found at end of
tunnel. & vericle
a paper or paste
marks & location of
web. & itch mite
entered & spider.
You will usually
find itch mite
at little distance from
this paper and
by means of a mag-
nifying glass show
power. You can
see & white at
end of tunnel &
can be removed
on point of needle
can be placed under
low power of microscope

E. Johnson.

Pinnis aeneus,
Hesperis.

two general
forms. (1) *Hesperis simplex*
(2) *Hesperis* ~~forma~~.

Hesperis simplex has
two classes. *Hesperis*
lobulosa or *fruticosa*, &
Hesperis prostrata or
Hesperis prostrata,
Hesperis ~~*fruticosa*~~
= *fruticosa* or *coll*
rosa. It is most.

only occurs about
1 inch out of water
It begins to hyperemic
base upon rock. and
red poplarly, appear
fruticosa like poplar
very round and

almost certainly
ecolies. In addition
to this you find
curious oritch
tunnel your diag.
is certain

Itching is very
marked especially
after & post. gets
in bed and gets
warmed up. These
post. claw themselves
& scratch. they cause
excoriation of skin
wh. becomes dry.
inf. giving rise to
a secondary dermatitis
wh. is ^{dermatitis} ~~itch~~ more
marked than that due
to & ecolies.

change in color
from white to gray

into vesicles, the
vesicles will vary
in size from a pin
head on up to a
split pea. When
they are thickly set
they are very coarse
giving rise to little
bubbles or vesicles
is at first clear
but soon becomes
cloudy and there
is considerable
tendency for these
vesicles to rupture
leaving a little
superficial ulcer
on their base.
Generally however
they will dry up

genital organs in
male & gluteal
region of both sexes,
especially about tubercles
in skin. It is also
reflected for along
while it will cover
almost entire body
except face & scalp.
In infants & face more
affected due to feet
& cheek rests against
body of mother when
they have scabies.
An itching dermatitis
of papules, vesicles
a pustular form that
shows a marked
preference for areas
just described &

without rupturing
of coracoid ligament
with 2 cords of 2
on 9 days to 10th.
they were dug up
into a scab. bled
freely of leaving a
hyperemia on the
skin drooping
in white while.
At these little
necroses get
deeper than can
in wch. even they
become rose and
these little ulcers
may persist for
considerable time
the lesions of fever
disturbance only

hatch & young
female porcupine
make new tunnels
in skin. The male
pinto does not tun-
nel a skin.

Sympt. of scabies,
that the will be
papules vesicles or
mice. pustules which
show a tendency to attack
& interdigital folds,
fing. & palms. Then
flexor of wrist,
elbow & axilla.
breast & female
especially around
nipple & ant. abd.
rashes in both sexes.

black & grey

2 or 3 or they may
be all around the
mouth & on
sides of nose.
But they appear
on cheek & on
forehead. They
seem to be more
common in women
and children than
in men. They fly
follow a natural
of typhoid fever &
pneumonia and
they fly appear
on lips & around
with disorders.

There is supposed
to be a nervous
element of block

~~the~~
burrows will be
from $\frac{1}{8}$ to $\frac{1}{4}$ in long,
at either end &
when it can be
seen it looks like
a ^{dark} ~~dark~~ coloured mark
just under epth,
is called a *curriculum*
or tunnel and its
presence is pathog-
nomic of scabies.
When you find it,
it is a white long
egg and burrows.
Along underneath
skin & deposits
excrement. These
eggs hatch out in
a course of about
10 days, & then they

of these changes
the charges
on an upper
layer of con-
crete - the charge
on those of upper
concrete with slight
round cells in-
filtrate & layer
of concrete. with
fluctuating overlying
surface. and a
venice.

Treat

Drinking pro-
tection. and you
get hold of a condition
right in beginning
of the of concrete of the

scaly. The remedies
should be applied
for considerable
time after the
appearance to be cured.

Lecture Nov. 3rd 08.
Scabies = Itch
Scabies is an animal
parasitic dis. charact.
by its papules, pustules,
thickened, & shows
a very marked tendency
to itch. Scabies is due
to the itch mite or
Acarus scabiei. Female
mite burrows in
epidermis, never going
into the true skin of
verte.

at the mouth of the river

ied locally may
of about this nature
on alcohol doubled
giving ^{great relief} quite relief
when the
comes, fully devel-
oped try to keep it,
from being ruptured
by advancing core.
When they do rupture
of ulcers from the
cores should be in-
cluded with a stick
of Agard's sticks
but avoid being
about rapid cure
then use some
single protective
ointment. Rules
find out or weak

~~of the face~~
of during your
treat. & after having
in place you
are required to
with the following
remedies, and
use soothing
lotions as. The
& Colamine & Berck
wash & Colamine
lotion

R. W. In fact it
difficult to cure
the hairs should
be removed either
by depilatory or by
pulling them
out by force.
You will cure
these cases as of

many things I found
much more to

carbide light
when they can
in the corner
of the room & give
use on the wall
it was kept up
definitely by two
men of light. These
could have been painted
over with the Ben
gon two or three
for boy. I Benjo
leaves a protective
layer of pretty
body of boys
point to collision
over the audit
will serve as a
better spirit.
have to me, one

adrenal medulla

It gives rise to
ring like areas
with inactive
margin and
clear towards
center. The fungus
gets into the
follicles and
than follicles
just like it
does in R.W.
It spore. A
further condition
develops on
face the neck
of the ovum
parts involved
A. B. C. similar to
that of R.W. of
nolp.

the most common
of venereal diseases

advised nichols
phosphorus & arsenic

James Fort.

~~James Fort~~, or
Vincennes.

W. J. to be ac. infl.

It is caused by a swell-

ing of the venous

system and hyperemia

of the affected area

and consequent un-

-der. H. Gibbard

previously more than

four in a affected

region & out of

four depends largely

on age of individual

in children and

adults up to age

— ed r o c h

v spread. the inf.
heads must be cov-
ered with cloth
age to prevent v
spread of v dis,
about puberty
it will get well,
diag. of R. W. of
v death. from
Ergasia, Schenckia
promosis.

Ring worm of
v Beard. =
tinea Barbae, =
tinea Sycosis.
this attacks v beard
& face seldom beyond
age of 50.
It begins like
R. W. & rarely does

I think the ink is
very light & muddy

of so & point me,
very light & muddy
to. I with life and
old age. & point me
great. Old people may
go to bed with their partners
frequently in a few
hours or perhaps
after first point
indicated little
hyperemic areas
whether they appear
on these areas
some very ~~many~~ small
poxes develop and
rapidly become vesicles
near these vesicles
also very high
poxes & in head
to erupt from.

W. L. ...
If you become
of support with
to know you will
be able to cure
your ~~for~~ R.W.
quicker than
by other ways
to know is applied
to & kept until
chain falls
then your for-
mation is applied
vigorously & by & time
2 mon. the ring will be changed
the return which is brought

In institutions
where there is a
great number
of children great
care must be
exercised to prevent

st. however they
are not longer than
anatomical seeds
when thickly set
they suggest the
to a very soft
line bulged or beeb.
these rolls of ves-
icles are usually
rather tough &
show very slight
tendency to split.
On protracted days on
ypts to these vesic-
les drying leav-
ing thin scale
like. some fall off.
As in fact they give
rise to no tearing.
But st. they are not

med after it
appears to be well
20. Crown oil
is used over &
affected area to
prevent any inflam
this inflam
destroys V.R.W.
finger but great
tense must be
exercised in &
used crown
oil of oil can
only be used over
a small patch at
one time in of safety
Kerion: does not
need anything but
nothing operation

you wrote to me

few days before
the shot for there
re-reveal crop of
the berries in
which went & it will
be prolonged for me.
to 2 or 3 wks. there
is only one patch
on the side - along
the spots - they
may occur over
distribution of
one ant. nerve
or they may occur
over 2 or 3 ant.
nerves & continue
to decidedly more
p. over & internal
region. It is
will be more

2/1, ~~some~~
should be removed
by a depilatory of
Rsa sulphuric &
ZnO & Cret. Prep.
For place over
it place and let
it remain until it
dries; then
apply a side
should be applied
that is mention-
ed under the
other forms.

As a hair regrowth
it should be removed
again. In time,
this R.W. fungus
can be easily destroy-
ed but the heat
must be contin-

the old covered
passage

instances $\frac{1}{2}$ or more
of the covered
path being
vehicles. This is
rare - the bed
occurs on only
one side & is in-
complete - There are
just enough cases
which occur on both
sides to prove the
rule. It is
It is a great over
distinction of any
entire. It is
infr. over a distri-
bution of 15th Ave
it keeps a relative
division and
it is a keeps

Transverse

V.R.W. may get
some into v. ruber
cut, this

V.R.W. figures can
be demonstrated
under a microscope
by extracting oil
from seed dropping a
few drops of 10% KOH
and pressing down
a cover glass and
examining preferably under
oil immersion

Trent's vegetation
requires a long time
to grow over 1000
acres should be clipped
- fed short and
then over V.R.W.
neighborhood

18.15
Jobbham 18.15

Zoster occurring
surface that when
other symptoms
for which develops
to have in 1 aged
it may come
by a secondary
infection going to the

Lecture Dec. 8th 08.
Herpes Zoster (cont'd)

Symptomatology. (cont'd)

In the neurotic group
but after 1 lesion
has disappeared. In
majority of instances
to 3. It is superficial
Does not give rise
to moving but it
1 lesion necessary

~~It is contagious~~
It is contagious
to me. gotten from
domestic pets
such as dogs
pigs and
also from cows.
It may attack
otherwise wise
healthy but it
shows a preference
for those whose
health is below
normal. In other
forms of R. W. the
fungus penetrates
the hair follicles
& grows and my-
celium and
fruiting
spores.

March 24, 1897

may even become
gang On either of
these went rearing
will remain. In
H. Z. of a few ulc.
is especially likely
to occur in aged
it may never fully
in very rare cases.
H. Z. is associated
with paronychia. That
is a paronychia it fol-
lows H. Z. - This is
very rare. In H. Z.
of a few you may
get paronychia of a
few nerve.
In herpes of occur-
ing about 1/2 and
about genital

have not made

pain is not
very much the
kind of R. W. is
spoke of as time
Kenton
when R. W. of
scaly gets up
which is repeated
except in the Ken-
ton where
loss is perma-
nent. Fortunately
by Kenton is
not very prom-
inent

R. W. of scaly +
a def. of child -
good as dattely
both sexes alone

January and

again you may
get paralyis of
upper & lower
limbs of winter.
Is very rare and
paralyis gets
better in short
while in nearly
all cases.

Cause of H. J.

Not well
understood. Is closed
among & aged -
nervous & is freq.
associated with exposure
to cold & may
follow some of the
reverse inf., such
as pa & typh - fever,
etc.

sets up a sharp
infl. and gives
rise to a condition
ash. looks like
Corymbosa.
The hair roots
fall out and
follicles open.
In some points
and they discharge
- a mass of pus.
The entire patch
border & center &
circumference
is involved in
this infl.
It causes no con-
stit. except I would
mentioning the

hyperemia

often between
ages of 10 to 30.
Is supposed to be
due in most
cases to overstimulation
of the vagus nerve & ganglion
or part of it. It is
very rare. In some
cases the vagus nerve is
not involved but
the nerve peripheral
to it is inflamed.
Partial changes in
the rate of
circulation.

Is that of
hyperemia with some
inflammation in
the vessels with ves-
icle formation on the

now - 
somewhere the
small patches
have ~~run~~ ^{run} together
together there is
a form of R.W. of
scaly, irregular, and
tufts of hair fall
looking & scales
shortly these tufts
will fall in
patches here and
there, this is
spoken of as dis-
seminated R.W.
It is common
in those whose
health is below
par that a R.W.
fungus penetrates
very deeply and

Ornithoglossa

ing in here.

Ding:

Ornithoglossa
very you may
find difference
from ~~scylla~~ red
but occurs on
you from ~~Hyper~~
Focies. ~~cells~~ from
Ingesta Contagiosa
web. occurs on

you. the Ding is
based on these points
in presence of hyper
-gum patches strikingly
net with vesicles

web very in size from
in health first few.
The. long when cool
even be unilateral

~~the~~
off and there
the stubs appear
as black dots
hence the name
black dot ring
worm. the loss of
a burr might be
quite extensive
you will find a
dis. more active
at a border

you will find
some greyish
white dryish
scales scattered

over a area

there are only
a few small. spots
has present on
there small. large

in distribution of
occasional
distribution of
sensory nerve, more
often intercostal but
may be elsewhere.
H. G. Is frequently associated
with neuralgia, pruritus,
the subjective symptoms
such as itching, burning,
stinging & some
tingling. The
subjective symptoms
are related
to the peripheral
P. n. s.

Do good. However
not good on those
who are very weak

It begins as a brownish scaling over on the scalp which spreads just as R.W. does elsewhere. In a short time R.W. penetrates the hair follicles & the hair itself. Being at great depths and for a long time it is so hard to eradicate. After while the hairs become brittle & they break off just above the level of the scalp at times a great many of the hairs for break

Of being a ...

You must be a
little bit con-
sidering of
become the
persistent
give looking some
days before
for a long while
after it has dis-
appeared. As
as a rule it runs
its course in a
week or 2 weeks.

It is unilateral &
it occurs only one
time. The person
is unimpaired, tho in
rare instances there
may be more attacks.

Jan 1

Sodumilyposulphate
is good and elegant.
You course of you
there. infl. appears
you stop your
forniticide and
apply nothing later

Timea tunicans =
Timea Copius =

R. W. of 2 scaly

R. W. of 2 scaly is a
dis. of refuted blood
very rarely develop-
ing after age of 15
and nearly always
subsiding about
1/2 age of steps puberty
even if kept
untreated.

Godwin's or ...
... the ...

That

is much
very simple. It
present in a spot.
whose general health
is down be
must have some
our great ~~importance~~
of ~~power~~ & ~~then~~ ^{recovery}
it self to relief
of pain.
General character
to ~~be~~ ^{be} ~~the~~ ^{the} ~~case~~ ^{case}
up for a pair
at ~~the~~ ^{the} ~~same~~ ^{same}
required. Phosphorus
of ~~the~~ ^{the} ~~husband~~ ^{husband} ~~or~~
bitter or ~~some~~
to use. As beginning
to be left out in

* Bichlor. 1-250
the most thing
is to be diligent to
your treatment and
you continue your
treat. for a consider-
able while after
dis. seems to be
killed or rather well.
Chrysorhin. in
Oint. 30 - 40 gr of
the oint. must
be rubbed ^{thoroughly}
about twice per
day after scales
have been ^{removed} ~~rubbed~~
with soap & water.
the lotion must
be put on two or
3 times per day.

even for some

there should be
applied locally with
vamp. sympt.
subside the
used some of
parasiticide.

Oint. contain
13 of neplan
to the 3 of paroline.

Beta-Naphthol 3T
Ammon. Hg. 3T
to the amount of
base then you
conclusion

Va. hyposulphite 3T
Agar 3T

more recent works
& vessels should
not be purchased
because if left
alone they dry up
in a few days. Just
using a sprayer than
a dip. Of course
to be avoided.

Iron in a solution
of white lead water
& borax applied as
a wet dressing
will give relief
usually than that
of any & rubbing &
dry dressing is
preferred. That is
ZnO, oil. or
borax oil.

the deeply outlined
of border, & features
the reading center
and the presence of
v. min. vol. are
present. Gravid
there is no more
typical Br. W. patches
near v. region.

Treat:

Is satisfactory
if there is much
infl. present you
will require to relieve
this by soothing
lotions such as Dead
water & lard
and Colamine
& gr. O. lotion.

the black & white
magnolia

Continuing to our
boat + 2 of course
assumed however
it was more
convenient to use
some of roosting
found much
of the structure of
the house was
on single beam
and there we
dusted over the
and then a protect-
ive dressing placed over
this and finished

Demonstrating Wey-
ipon.
Is now.

magnolia up of you^{like}

lge. patch. It
It is spread
onto & gluteal region
onto & pubis. or onto
old, wall and
onto & axilla.

Its extension is
quite rapid and it
is wide spread.
Its extension is
usually confined
to & crural region
there is often a good
deal of itching
which is very
intense and as
marked as that of
eczema.

Sitz: Usually
fairly easy

~~The world is~~

Pumpkins

of course

~~curious to know~~

myths in which bees
own pumpkins but
you get bees in
apple & depository
in Argentina & in
lungs. Bees don't
these causes don't come
under clove cases

Pumpkins
Pumpkins in
count - dis. choroid
by development of
a greater number
of bees more or less
symmetrically
distributed upon
hops better still

Winters line
in the face

brownish scaly
patches as R.W. of
V body there is
not on one
patch there
spread from a periph-
ery. Is more
markedly inflam-
matory than R.W.
of V body, so that
of advancing margin
is higher raised &
redder. at times
there is a good deal
of clearing in
center practically
always or deep decline
this form of R.W. me.
confined to only
a few patches or one

from January

distance from
united. ~~the~~

courses and areas
about the

there are 4 ^{sub}divisions

1. ch. Purple.

2. ch Purple.

3. Purple - bluish

4. ch Purple - Vegetation

ch. Purple

to by for

most a society

Things are. pe-

ceded by count - right

which are right. Ah &

please with me

probably with they

me about a 20 f

wild and ^{attract} ~~attract~~

no attention

On you come
in hypomelence of
with T₃ & c. J.
Jword.

Lecture Oct. 27th 08.
Tinea Cruris. =
Eczema marginatum.
It is from R. W. Welch
usually attacks &
scrotal angle,
It usually extends
from scrotum
onto adjacent
leg. At times only
scrotum are in-
volved but as a rule
both leg & scrotum
are involved
It begins in rule

✓ bright begins to
bleeds well - reach there
full development
in few hours to a
day or two and
begin to rise from
capitulum to other
egg - in 2 days they appear
to move from the
skin there is
unreigning inflam.
violence and then
it is rise, present
that there are only
a few blebs on the
in a good many
neonly sloughs by the
- need to both rise
up body & they
more they over on

with some more

me. used.

Doline pitted over
a R.W. will cause
✓ R.W. to happen
after several
applications.

Solizic acid
may be used.

20 grs to 1/2 of V
announced Hg
me. used, it
allate of Hg me.
used.

Rx.

Solizic

Heplm

Baseline

XX grs 5 1/3

3

1/2

3

1

our little copper
portion of the
stick is a piece
they may own
any where else, now
of not any more
the trace of the stick
is gone and the
contents are all
at first, room
became cloudy.

and then more
as close there
we saw beds

formed out in
the bed. 1 individual
leaving usually
nubrids in 10
to 10 days. the con-
tents of the bed.

Health is poor
from any cause
than those whose
health is good.
It will fly, run
its course and
disappear without
treat. but as a
rule some of the
simplest parbit
inides is requir
ed to affect a
cure. In the
more inflan. cases
of R.W. nothing
application such
as Leadwater &
Santonium and
Zinc oxide and also
Juniere lotion

dy up at p
a thin scale wh.
won free of pro-
ing appearance of
with good return to
normal color.

Successive crops
occur from time
to time generally
~~to~~ without a rest
is not needed over
considerable
time ^{months} ~~the~~ ~~recovery~~
Recovery is rare
in cases of panga ex-
cept ch. panga
the recurrence
are not rare
& cause die of
exhaust.

~~the patches~~
A R.W. patch will
ply reach a diam
of 1-2 or 3 in
of which there
are several
patches together
they may coalesce
at the line of
coalescence being
seen and give rise
to a larger more
irregular patch.
This condition
may occur at
any age. I have
seen it in children
It may spread from
v skin of the human
It attacks
those whose

bee people,
 common right,
 removed
 accelerated fever
 accelerated pulse
 & bee-people food.
 upon occasion
 in a chronic condition
 & contents of blood
 & bee-people often
 become purulent,
 & reason for
 believing what is
 called bee-people
 is necessary and
 people but is
 pyrogenic itself.
 That honey is very
 rich in butters
 & those who are

of being ~~the~~ more
or less covered with
poplars & willows &
W. Juniper. This
rocky & R. W.
Juniper has penetrated
much deeper than it
usually does. In
times of rain &
R. W. Juniper is
curly & curled &
& Epidemic St.
It penetrates into
the and St. into
& cuts over and
in such events it
gives rise to
more inflammation
cases of R. W.

continued to
fumbling dead
meat with fly
terminator for
of course. is not
long and may
be proven

Pungia foliosa
is more dis. and
it higher as black
with may look
like 1 black of the
pungia. but now
fly & use is
not true but
is Cox. & contents
in black soon to
become pungent
& black wall breaks

1 portion within 1
 circle will grad.
 return to normal
 at 1 0 but as you
 leave 1 center of
 1 condition will
 become more
 marked as you
 reach 1 periph.
 1 color is usually
 light brownish
 1 scales are
 present in varying
 degrees. In some
 cases 1 center does
 not clear up, and
 border is very decide-
 dly elevated. Being
 somewhat more elevated
 than 1 0 and where

before long leav-
ing a row of retic-
les, and a pair of
long of red. long
adrenal epidemics &
beneath these top
process continues
to burrow extend-
ing peripherally
there is a space of
at least 1/2 to 1
a new blood on
a dyed site of
old one but there
being an epidemics
to cover it we
singer have a
oogy of. from
long cold. mi-
trick of point

Stomach less than

sch. is usually
slightly elevated
and sharply
marked from the
healthy tissue. ~~But~~
In films. Under
is slightly elevated
at other times
it is sharply elevated
and contains papules
and vesicles and
showing. Consider-
able infl. reaction.
In most cases
however elevation
is very slight.
Papules are very
poorly developed
and covered with
scales

Wash. & ...
New York ...

Of these birds we get
thickening of skin
epidermis ^{exposed} & ^{exposed}
flesh of ^{exposed} ^{exposed}
~~the~~ there joints
are especially
likely to occur. Great
distances of joints
to the whole
then - birds &
beats. It becomes
wide spread in time
to you get large
patches of this thick
and ^{is} the web.
is constantly re-
creating and
and thus dipping into
cuts. Cont.
myself - one point
usually goes for

Thin Circumference

is 1 more an

form of R. W.

It simply attacks
1 neck and arms

fore and hands

It may appear

on any of them

hairs of top body

It begins as a slight

elevated papule with

the grad. spreads

peripherally. and

as it grows

older & longer it

begins to clear in

center

1 active portion

of 1 R. W. is at

this border

Grade ~~note~~ ~~note~~

One in a quart hill
these for get let-
ter as a nice try
are from ~~exhaustion~~
the ~~in~~ ~~an~~
they attacked much
our mouth & throat
to put ~~cost~~ ~~cost~~
mellow & well,

4. Pangha. Vegetation
begin the pangha
piliocoon when
bleed bleed well
my ~~first~~ leaves
are ~~secret~~ ~~secret~~ ~~secret~~
of ~~become~~ ~~covered~~
with vegetation
due to ~~the~~ ~~presence~~ ~~of~~
the ~~dense~~ ~~populus~~

~~the~~ may attack any
portion of body
but ~~may~~ ^{often} occurs
about ~~menstruation~~ ^{menstruation}
summery hills.

Jan. 5th 08.
Etiology of Pimple.
Not well understood.
Seems to occur at
pretty much all
ages. Perhaps more
frequent in female than
male. Any condition
that depresses & gen-
eral health seems
to render one more
subject to pimple.
Is frequently known to
follow sudden chill-
ing.

96 Mch. Johnson Chase

or train from 1 follicle
and then 1 paras-
itism can be
applied.

This treatment can
be repeated and
continued on for
a long time even
after the traces

of it have
apparently dis-
appeared there
is likely to be a few
of the germs left
behind it is
likely to reappear

P

1. I have been 1/2 please
2. I have been 1/2 please

25 July follows those
who have been under
a great stress for a
time, as business
worries etc.

26 more p. in ner-
vous disappointed
people, and it follows
nervous frights. It
is July, and it is with
out any dis. of ner-
vous system & the
more July it has
been associated with
functional nervous
trouble. It is prob-
able that people is
dependent upon
nervous conditions
than any other.

Just finished with

to be some other
a thing though
and with that to
join apart.
pull this part
vigorously and
dis. arch and al-
low it to remain
until it begins
to sweat. Work it
off and it will
take & burn the
longer it can be
allowed to remain
& deeper it will
follow & burn into
the follicle and then
can be repeated
as long as is neces-
sary to get relief

the same way
knowing that

Just. oriented with
gastro int. in to-
tion and after
mins. have been
brought forward
to explain & course
ac. periph. is almost
entirely due to some
of the pyogenic mins.
of path. of periph.
It usually comes
on unassociated
with symptoms of infl.
or they are so slight
or so almost escape
notice. Has usually
more or less sudden
develop. of swell-
ing in region between
of 2 layers of peritoneum.

15. at present between

Use, Sulphur
Sulphur. $\frac{3}{55}$
Calcic chloride $\frac{48}{4}$
Base line $\frac{3}{1}$

Or you can use
calcic acid and
announced by
Or Colonel, old
Hq. I put any
or paratinder, will
be the work

In children this
pulling of a hair
is rather different
and objectionable
so you may use
a depilatory.

Barium sulphide $\frac{3}{111}$
Zinc oxide
Cretan prep. aa. $\frac{3}{1158}$

109
St. it occurs between
basal layer of rete
& papillary layer
of derma. In pemphig
regions there is a
rolled up part of
the papillary layer
of the vera
Diag.

Usually
fairly large.
The most difficult
dis. to diag. from is
dermatitis herpetiformis
if you are not a
skin dis. In derma-
tis herpetiformis
the dis. is more widely
distributed and
the lesions are or

~~nothing to report~~

be clipped very
closely or better
still pulled out
over affected
area and for about
 $\frac{1}{4}$ in all around
and then par-
nitride applied,
a good one is
Hyponitride of
Soda 13 to 13.5
H₂O, vigorously
rubbed in twice
per day or less
keeping a constant
motion to application
to the patch.
Sulphur and
sorbic acid oint-
ment is good

roughed in a near
fayer and we have
intense itching
which is not present
in purple
Dingertgo Contogiosa
is a dis, most freq.
mistaken that is
below for
distribution of
impetigo Contogiosa
is diff from pyzsh
being on the hands
& face etc. parts
reached by l. hds.
the ac. course &
evidently contog.
nature of impetigo
contogiosa. These
should make a diff

James G. Thompson

One of the best
methods of treat-
+ quickest method
of cure is by the
application of a
gray mixture
to make a
fool and when
it falls & vigorous
use of parasiticide
until it effects a
cure, then I move
on less a longer
cure. Because too
much demerol
use, prod by
use of X-ray & has not return.
When X-ray is
not used & then
should after

consequently easy
H.V. belongs to
A. unicolor, leprosa,
I. p. & any length
or color with bees.
P. unicolor is
often mistaken for
unicolor, but
unicolor has other
symptoms of unicolor
which render it
highly likely easy.
I. p. & unicolor
helps in diag.

~~Dna~~

Prozi

In all forms.
it depends more or
less upon extent
of range. I. p. more

or be detected the

it is hard to get
a parasitised
in contact with
it. This mis-
conception can be detected
by soaking some
of scales in 10%
KOH. and then
staining with
some of Laitine
dyes.

Prog: can be readily
located in a short
while when it
attacks & body
of it can be
checked where it
attacks & scales
by persistent
test for a long time

or less extensive the
prog. ~~at~~ grove
of not so much
not so grove,
death in

See V chronic form
V prog. is usually
good V dis. may
run for wks. or
mon. or longer
recurrences are
almost certain.

It ply happens that
V ch. form of pemph.
wears V pat. out and
kills them. pemph.
infection is much
more serious. It
may last for several

From back of a mule
seen a very small

which is a vegetab-
le parasite that
grows by propa-
gation in the hair
substance and
hair follicle.
This disease is a
very ch. cause
it will continue
on the scalp until
indefinitely until
the hair is destroyed.
Where it occurs on
the gen. body of it
is rather easy to
cure. Where it
occurs on the scalp
it is very difficult
to cure because it
gets down so deep

years but as a rule
in the course of a
few months or a
year or so it tempo-
rarily settles itself,
however in the
minority of cases
it does not recover.
Single vegetation is
almost always
fatal & kills within
3-6 weeks.

Treat

1st gen. health
should be built up.
Nutritious food and
general tonics are
indicated. Arsenic
quinine & strychnine
are especially

or has any
in eastern
cities.

Both eyes are
equally affected.
I know of
no one else than
in those who
are couple. Do
more common
those who have
poor health. It is
common, but not
so abundantly so,
as in those who
where there is
crowding. Do due
to Acetone
Schoultz

no more work

indicated
As. has enjoyed the
rep. of being a
specific & natural
ly given as forcers
soft beginning with
usual. Does it run.
& good I until
you get v. phys. effect
then decreasing
your horse to the
point at which v
not even standing
for a good long while
As. is by far v. most
simple. But remedy
for pump.
When bleed are
present in v. bullous
forms therefore

Ground Losses

then into v structure
of v hair follicles
v ch. infl. ^{Changes} ~~changes~~
causes v loss of v
hair and atrophy
scarring.

Favus always begins
in children. It ~~very~~
seldom begins after
v age of 15. But in
these cases it may
last well almost
adult life ^{even}
into old age.

Favus is not very
common in this country
except in newly
arrived immigrants
and so makes its
appearance in

the same as in the case of the
other two

usually found.
under aseptic pre-
cautions. there
epidemic covering
allowed to remain
or dressing,
where the blood must
be removed. becomes
now some soothing
lotion such as the
Calamine & Quinine
or carbolic oil
make the best dress-
ing in the follow-
ing variety. the con-
tent of both is jelly,
employed then
under aseptic slightly
antiseptic dressing.

Mr. J. M. Smith

until it destroys
the hair and then
it will usually
get well. Fungus
may attack the general
body of a hair and it
forms more or
less circumscribed
patches and may
be widely spread
the superficial layers
are developed but
are not as well
defined as when
occurring on the
scalp. Fungus may
penetrate the hair
follicle and then it
finds its way into
the hair shaft and

make the leaves
change in color

you may use (qu)
+ colored soap,
on colorized oil.
In people's regions
your look is hope-
less but supportive
that cleanliness
of washing applica-
tion will be the great-
ly to your comfort.

Demat. Herpeti-
formis. Skipped.
usually fatal. For
Droon.

Imperio Herpetiformis
skipped.

P. varians.

Isach. info
Dis. of skin, Character

12th May 1910 - 12th May 1910

white lose their
sneplum color due
to accumulation
of ordinary scales &
dirt. In time
they will come &
begin to fall out &
this hair loss is
permanent and
causes an atrophic
condition of the scalp
and then will be
a some operation
beneath the crust
due to irritation of
scalp by their presen-
ce. X-rays shows
but little tendency
to spontaneous
improvement

by develop. of properly
defined patches of
the elevated base.
slightly above U
then covered
by scales wch. are
characteristic. Proliferation
occurs at all ages
but usually not
before age of 8 or
after 40. It begins
as a slight papule
on apex of web, is
a silvery colored
scale. this papule
spreads peripherally
to form as it remains
a papule it is called
Proliferous punctate
the this papule grows

Hydrolytic skin - 8

yellowish crust
pierced in center
by a hair this
crust will 2 in
size and will grad-
ually become lens or
saucer shaped.

Its convex. γ presses
against γ scales
& causes some atrophy
of γ skin. some
depression. There
are usually a good
number of pustules
and in badly neg-
lected cases they
fuse together and
form mouton like
masses.
and they will after

yellowish brown
lenses - 1 corner

2. - size by peripheral
extension, when
it reaches the size
of a drop of water is
= *Psorionis guttata*
then as this drops
in size reaching
size of coin = *Psorionis*
marginata, as the
psorionis patches
get older they clear
up in center &
= *psorionis circin*
ata - fully reversed
here with one side
to gether & coherence &
we speak of that as
psorionis marginalis.
V Patch of *psorionis* is
very sharply outlined

myth of some sort

~~2nd~~ 2nd Lecture.

Lecture Oct. 13th 08.

Dermatology.

Trachoma, Trachoma
is a vegetable para-
sitic dis. Character. by
formation of deep
cup like crust
wch. occur most

frequently on scalp.
It forms a mass of body,
as a rule however
it is confined to scalp.
And it 1st
begins as a little rough-
ening of a small
area wch. is covered
with ordinary scales

In a short while
there will be little

2nd specimen.
Specimen 105/13/18

from mura skin
its dull red
color slightly red
~~not~~ elevated.
Is covered by silvery
scales of chord
color, the pearly
on silvery. These
scales vary in quan-
tity & size. They
are numerous &
overlap like shingles.
At times they are
very abundant and
stack up on one on
other & one con-
stantly being shed
so that a typi-
cal & marked core
the bed shows between

when the pot-keeps
on are covered with
scales. If the coes
scrape off the pots
you will find little
punctate points of
pustules. It is due to
fact that the papil-
lae or papillary layer
of skin is exposed
at those points. As
the patch grows older
it tends to clear
in center & is
quite devoid of
protrusions of con-
siderable duration
several of these may
run together giving
rise to a large

patch of wavy and
line - It has quite
general distribution
generally first about
extension of sp. of
elbow thick.
as well as noted
with there you
will find patches
in scalp especially
of about 1/2 inch
head and monoid
process. 1/2 inch as
as well as scope, but
in extensive cores
1/2 inch also is im-
bored. 1/2 extension
sp. back especially
about 1/2 inch
then in extensive

covers the back
✓ rolls off feet & pollen
of hds. may be also
involved. From
find lesions in
the hair & elbows
etc. - I noticed a
pore in ^{winter} ~~the~~ ~~leg~~
It fly disappears in
the summer
itching may be
troublesome in
proportion of cows &
gives them no
trouble whatever
of depurating.
General health is
generally & exceedingly
good. General
and count to

plaint. Is very
chronic. and is
subject to recur-
ces

Lect. Jan. 12th 09,
Several varieties of
Pruritis. Have been
given.

Etiology of Pruritis
not understood.

Has been spoken of as
dis. of good health.

It occurs in both
sexes pretty much
alike and is more
inclined to occur
in those whose
social & other
duties are heavy

Very seldom
develops in young
seldom appears for
1st time after 40.
It appears during
preg. in female
occasionally it
has been known
to develop after
slight trauma
as a rule v. pot. me.
well but it is associated
with dis. that run
down v. consist.
some dispute as to
pathol. (exposed) of
proliferation. As usually
described as being a
hyperkeratosis.
Associated with a

marked I. of v rete
cells. v rete papillary
projection of rete
become consid-
erably enlarged.
v layers of cells above
v rete membrane
fall off before their
time. Hence that
way the papilla (open)
is exposed. v gray
color of scales, is
supposed to be due
to layers of air get-
ting in between
v cells. there is
some hyperemia
of papillary layer
of skin & is found dilated
and there is

some infiltration
Diagn:

usually
easy. It depends
upon location of
lesion. Upon
sharply circumscrib-
ed lesion. Upon
color of bone. Upon
character of scales.
Upon prod. of
little bleeding points
when scales are scrap-
ed off. You may have
to dif. between psoriasis
& squamous eczema
Between psoriasis &
lym. erythematous
Between psoriasis
& tuberculo squa

on sph.
reborhoca & Ecyra
reborhocaum.
In Ecyra & scales
foundations not
so great - scales
and not silver
white as in prothesis

In prothesis
unusually rocks the
cheeks.

Ecyra reborhocaum
& prothesis are. into
taken, scales are
gray.

Prog. of Prothesis.

As for an individ. lesion
is concerned & prognosis
good there lesions yield
fairly readily to treat
in ~~the~~ ^{the} ~~lesions~~ ^{lesions} ~~proceed~~
but recurrence is a
rule and permanent
cure can seldom be
promised. Treat. is
both general & local.
If for any reason a gen.
health is below par.
it should be attended
to. As. is supposed
to have some spe-
cific action. Every
popular ~~in~~ ⁱⁿ ~~treat.~~ ^{treat.}
previous & still
has ~~reputation~~ ^{reputation}.
And as follows

sol giving 1-3 gr.
after meals for
some time. Should
not be given for too
long at time because
it may prod, discolor
stomach & keratosis.

As. can be given in
fairly large doses
if K.I. have been re-
commended, 5-10-
15 gr. tid. is felt
in the very large doses
are given. Crocker
reports excellent
results from Solcin
in about 15 gr, doses
tid. & use of alcohol
has also been favorably
reported on.

Liquor Potassii is a
favorite one.

Best remedy one
gotten from local
& combined treat.
Chrysarobin is the
best, perhaps. It
has to be used with
care because of its
tendency to prod. a
dermatitis & if used
about V hdt and
face it often prod.
a rather sharp con-
junctivitis. I used
it then in V form
of oint. in which
we use 10-15 gr
to $\frac{1}{3}$ of oint. base.
You may use it

~~with~~ solingie and
10-20 gr. of $\frac{1}{2}$ z.
+ resorcin. The
resorcin. permanent
ly stains underneath
the paper. I only give
with collodion.

Rx: Resorcin gr 30
Solingie and gr 30
Flexible Collodion $\frac{1}{2}$ z
Wash in paint on the
lesion with combs
hair to brush 2 or 3
times per. wk. Oint.
to be used 2 or 3 times
per. day.

You may use trans-
mucin = liquid
butyrola instead
of collodion.

Before applying
liquid remedy
must remove
moles first & give
room both with
some soda in
water - If it
very wide spread you
had better use this
chrysomelid over
a very sharp and
a good sized
general denudation.
On this patch
it could under
influence of chrysa
robin it becomes
white white &
surrounding skin
is yellow.

✓ tor. prep. are
used. Either ✓
ordinary tor. or
oil of fusel oil
of cold. are used
from $\frac{1}{2}$ gr up to
2 gr to oz. the ob-
jection to them is ✓
odor & color. They
can be used either
in oil or in col-
-lodion. ~~✓ HgCl~~
✓ mercurate prep.
are good especially
white ppt. can
be used in $\frac{1}{2}$ to $\frac{1}{3}$ to
3 gr in oil.
Sulphur me. and
sulphuric & oxylic
acid me. used to

after in fact.
Polycyst should
not be used about
1/2 free because the
conjunction is
liable to occur.
Provisionally, if it
appears in summer
& relapses in winter
when all other
treat. have failed
1/2 use of Kroy
for 4 or 5 min. three
times per wk. may
be used after treat
with Kroy. it does
not relapse so
suddenly. It is
harmless & must be
used carefully.

Imperigo con-
toziona

As water can dis-
solve & due to its form-
ing mis. little
grains small. blisters
in blebs surrounded
by an inflow. on the
the & blebs is very super-
ficial & its contents
very rhy. become
pusulent. They soon
dry & leave a thin
yellowish scale cov-
ering the bone. As very
lightly attached to the
bone & fly, ripped
up around the periph-
ery description of its
attachment to bone
is void & the kind

of stuck on. In a
few days the crust
falls off leaving a
puffy ~~surface~~ blue
web room returns
to normal giving
rise to no morning
cumber these
to vericle. may
conclude giving
rise to sleep

of some size. On
original vericle
may begin to dry
in a center where
it spreads purple
giving rise to a
vericle some
size. The individual
portion shows its

course in 4 or 5
days & I think
necessary crops
are in rule.

Goodly, almost
perfect, & chief
books but they
often & I don't
know in in
& corners & dirty
that they are
confined to that
clock. Distribution
is chosen by D. Am.
very begin around
& corners of mouth
those & heads
to other portion
of free nearly on
hand of fingers

Boole's Law

See first lecture
in the 1st Aug.
Olin. Boole,

See first lecture
the 25th

Very often present
garbles & duct
lice. It may attack
any portion of body
you find the great
majority of cases
confined to hands
feet & more
gives rise to itching
night & foot but
not enough to
ant. to much.

Is due to ordinary
pus this they get
entrance to thru a
slight abrasion. I
suspect it is due
to superficial layers of
epidermis & give rise
to no scoring

this condition is highly
contagious will pass
thru a family or commu-
-nity. The treat. is very
satisfactory. Must first
be soaked off with oil.
Iodine & mild
disinfective remedy used.
Potassium bich 5-10 grs
to $\frac{3}{4}$ Anhydrous
H₂O 5-10 grs to oz.
Keep scales soaked off
Wynne paper a little
wet. Boric acid is good
N₂O and H₂O & soly-
lin acid are entirely
sufficient

[REDACTED]

[REDACTED]

Dr. Knott recommends Stanton's
Blood pressure instrument,
for clinical work.
Ehrlanger's Blood pressure
instrument is the best.

