


SL 2 RCM PV
 -3.4 %



SL 3 RCM PV
 -13 %



SL 4 RCM PV
 -6.8 % -16 %



D

M 171

HR's INFLT - TENDENCY TOWARD ↓ DURING REST & RECOVERY STAGES
 MIN. VOL. " " " " ↑ " "
 DIAST. BP. " " " ↓ DURING EXERCISE

ALL COMPARED TO PREFLT VALUES. ALL INCONSTANT

VITAL CAPACITY - ↓ 2-10% INFLT. - NORMAL POST FLT.
 (SPT 1 SL-4 RET. TO PREFLT. NORMAL AT END OF FLT. - NOT OTHERS)

POST FLT. ↓ EXERCISE RESPONSE - MANIFESTED BY

$\left. \begin{array}{l} \text{BUT 0 IN 2/3 SL-4} \rightarrow \downarrow \text{C.O. (Q)} \\ \text{(45\% BUT 0 IN 2/3 SL-4)} \rightarrow \downarrow \text{STROKE VOL.} \\ \text{ALL} \uparrow \text{H.R.} \end{array} \right\} \text{FOR A GIVEN } \dot{V}O_2 \text{ (O}_2 \text{ CONSUMPTION)}$
 (MIN. VOL.)

ALSO ↑ RESTING VENTILLATIONS & ↑ EXERCISE (3rd and) VENTILLAT. IN SOME (4/9)

SL-4 (ONLY) MAX. EXERCISE IN FLT, ~~PROB.~~ O₂ CONSUMPTION > PREFLT.
 (TOTAL WORK PREFLT. IN INFLT. NOT THIN)

~~VITAL CAP.~~

RRT. OF EXERCISE RESPONSE TO PREFLT. VALUES:

SL 1/2 - 18-21 DAYS
 SL 3 - 5 DAYS
 SL 4 - 4 DAYS
 PROB. RELATED TO INFLT. PERIODAL EXERCISE

PERSONAL EXERCISE: - ERGOMETER ONLY

SL 1/2 AVE 31 WATT-MIN/Kg BODY WT.
 SL 3 " 65 " (107% > SL 1/2)
 SL 4 " 71 " (130% > SL 1/2)
 (- ± 10% > SL 3)
 + MINIKYM & METAL SPRINGS - SL-3
 + " " + TREADMILL - SL-4

M 71 - Ca Loss 115 mg/day > Preflt. → 3.5-4 gm/Mo. loss UNABATED THROUGH FLT.

P Loss 32% > Preflt. (24, 42 & 32%, resp., in each astronaut)

N₂ Loss 3.5 gm/day > Preflt. - (also looks unabated)

Ca Loss = .35-.4 % / Mo.

Ave. for 3 flights ± 150 mg of Ca balance Not

All despite Hi-Exercise Levels
 St. Neg Na & K Balances

(2)

M078 - Vogel & Ray Absorption

Calc

SL-3 SPT - - 7.4 %

SL-4 PRT - - 7.9 %

SPT - - 4.5 %

All others Neg

Rubens & Ulman - No Lines

M073 LEACH

BL. PRESERV. - No 120TA

CHEM. ANAL. BY { RADIATION
FLUOROMETRIC
~~ANALYSIS~~ GAS CHROMATOGRAPHY

STATISTICAL ANALYSIS BY { COVARIANT ANALYSIS
PAIRED T TEST

CHANGES (PLASMA & URINE) ALL $\geq$ IN CLASSIC. KORTS.

SERUM & PLASMA CH:

↑ INFL & IMMED. POST. FLT

Ca
P
K - v. se.
Creatinine (Fol.)
X Cortisol
Angiotensin I - ~~imm~~
Aldosterone

POST FLT P

CO₂
PROT.
TSH
T₄
ALDO

↓ INFL & IMMED. POST. FLT

Na
Cl.
Osmol.
X ACTH

↓ INFL & ↑ POST FLT

Glucose
Insulin

↓ Post FLT

cholesterol
Uric Acid
Mg
Coarct. Polyhydramnios
(CPH)
Bilir

UNIVES:

INFL P

Na, K, Ca, P, N
Aldo
Cortisol
17 Ketos

BOT VARIABLE

OH PROLINE
OH LYSINE

POST FLT

Epi.
Norepi.
Aldo
Cortisol

IMMED POST FLT
OH PROLINE
OH LYSINE

INFL 17 OHC's NO CH.

INFL ↓ - (THRODS)

ADH, Epi, Norepi, Uric Ac.

POST FLT ↓

Na, Cl, K, P, Mg, Uric Ac.
ADH & 17 Ketos

(3)

M073 (CONT.) - LEACH

FLUID & ELECT. BALANCE

No H₂O DIURESIS 15 (DAYS) - URINE OSMOL. > PLASMA
∴ ↑ SOLUTE EXCRETION

APR 11 (BUT 15 DAY ANALYSIS COULD NOT BE
DONE - REFRIGERATION ~~FAULT~~ LACK IN CM)

POST FLT. - TBW ↓ 1.7%
ECF ↓ 1.9% } BUT THESE ↑/WT. BASIS.

EXCH. K (⁴²K & ⁴³K) { SL2 - -7.7%
SL3 - -3.4%
SL4 - -8.2%
MIN. - 1.1 SL3 SPT
MAX. - 12.3 SL4 PCT

NONE OF ABOVE RELATED TO FLT. DURATION

M113 BODY FLUID & RCM LOSSES

SL-4 } 7% RCM LOSSES (MEAN)
16% PV LOSSES (MEAN) - SL-3 = -13.7%
SL-2 = -3.4%
SL-2 = -13.7%
SL-3 = -3.4%
SL-4 = -6.8%

SL-2 - 14.3% AVE
SL-3 - 12.2%
SL-4 - 6.8%
SL2 RCM ↓ 15% (MEAN)
SL3 " " 11% (MEAN)
FROM MEMORY
PV LOSSES ?? ± 10% I believe

RADIATION

MEAN TOTALS (RCM):

	SKIN	LENS	B.R. FORM.
SL-2	3 1/2	2 1/2	1 1/2
SL-3	8	6 1/2	4
SL-4	14-17	9-12	7

(> .1 REM/DAY)

CAREER DOSE ALLOWABLE

SKIN	LENS	BFO
1200	600	400

(4)

TOXICOLOGY

± 300 Compds ^{found} ~~identified~~ in OWS Atmosphere
(107 identified)

SL-2 not entered suit after series of plunges
due to probable (G/B tests) build up of
Tot. Diisocyanate + CO₂ = heating of urethane
foam

All materials screened carefully - esp. for CO outgas
& testing of fluorinated polymers.

Non-metallic mats for CO₂ & organics

Levels of acceptance established based on
ECS capability, check rate vs. outgas rate.

Pyrolysis products also (not toxicity)

Loss of Δ shield $\rightarrow$ Skin temp of 350° F outside
160° F internally

MICROBIAL

OWS AIR - SL-2 LOW 100 (END OF MISSION)

SL-3 LOWER 75

SL-4 HIGH 350 $\leftarrow$ COLONY FORMING UNITS/m³ air

PUZ ALMOST ENTIRELY TO TERRATIA MARCESCENS
RECOVERED ALL 3 SL-4 CREW POST FLT (NOT PREFLT.)
- PERSISTED IN FLT THRU POST FLT
QUARANTINE (IN NATAL PASSAGES)

OWS SPUR SITES

SL-2 $\downarrow$ AEROBES ($\leftarrow$ Δ STRESS?)
 $\uparrow$ (10X) ANAEROBES

ENVIRON. STAYED CLEAN
HOWEVER (conclusion of
analysis)

SL-3 $\uparrow$ AEROBES + ANAEROBES BY 1-2 LOG UNITS ($\leftarrow$ DURATION OF
FLT.)

SL-4 AEROBES CONSTANT } AS MISSION PROCEEDED
ANAEROBES $\downarrow$

$\leftarrow$ LOSS OF PROPIONIBACTERIUM ACNES
ALSO $\downarrow$ Δ SKIN OF CREW (RESERVOIR)

CM'S showed No Variation PRE + POST FLT. - THO DIFF. Δ EACH
OTHER (OWS CHANGES NOT REFLECTED IN CM'S)

FUNGUS 5 OWS SPUR SITES - LOW UNTIL SL-4

$\leftarrow$ MILDREW GROWTH IN LIQ. COOLED GARMENTS
(SAME IN AIR) (SAME ON CREW SKIN)

EVEN UNUSED LCC'S MILDREW IN CONTAINERS (RETURNED)
FOR SPREAD

CREW - AEROBIC - 17 MKD. POST FLT. ANALYSES - OFTEN $\uparrow$ BEYOND PREFLT
WHEN 30, PUZ TO $\uparrow$ NO. 5 BACTY, BUT $\downarrow$ IN TYPES (GENERA)
ANAEROBIC VARIATIONS NOT OBSERVATION ON MISSION DURATION (API 14 + SL-4 ARE $\approx$)
- FEWER NOS. + GENERA POST FLT. (EXCEPTION - SOME SPL AREAS SL-3) - P. ACNES

BODY MASS LOSSES (LAUNCH TO R+0) + INFLT. CURVES

SL-2 CDR $136\frac{1}{2} - 132\frac{1}{2} = 4\#$

SPT $170\frac{1}{2} - 163\frac{1}{2} = 7\#$

PLT $175\frac{1}{2} - 167\frac{1}{2} = 8\#$

SL-3

CDR $151\frac{1}{2} - 142\frac{1}{2} = 9\#$

SPT $136\frac{1}{2} - 128\frac{1}{2} = 8\#$

PLT $195 - 185\frac{1}{2} = 9\frac{1}{2}\#$

SL-4 CDR $150 - 149\frac{1}{2} = \frac{1}{2}\#$

SPT $157 - 154 = 3\#$

PLT $149 - 146 = 3\#$

AVERAGES: SL-2 = $6\frac{1}{3}\#$ LOSS

SL-3 = $9\#$ LOSS

SL-4 = $2\#$ LOSS (MEAN = CLOSER TO $3\#$)

SL-3 SHOWED GREATEST LOSS (59 DAYS)

SL-4 " LEAST " (84 DAYS)

MAX. LOSS (SL-3 PLT) - $9\frac{1}{2}\#$

MIN. LOSS (SL-4 CDR) - $\frac{1}{2}\#$

MICROBIOLOG. (CONT.)

CONCLUSIONS

NUMERICAL CHANGES IN AUTOFLORA NOT CORRELATED EMISSION
TOTAL NOS. TEND TO ↑ FOR AEROBES, ↓ T-R ANAEROBES DURATION
NO. OF GENERA & SPECIES CHANGES LITTLE FOR AEROBES, DECREASES
FOR ANAEROBES

YEASTS & FUNGI

BUT REL. ↑ IN C. ALBICANS

↓ POST FLT. IN AP; ALSO IN LAB. - EXCEPTION SL-3 (NOT
+ SL-4 DECREASED)


POST FLT. MICROB. FLORA - NO RUINING

(ROSS CONTAM. - NO (SL-2 PLT'S STAIN AUREUS NOT OTHER CREW MEMBERS)

YES (SL-3 SIT GOT ROR'S S. AUREUS PHAGE TYPE 3A
BUT NOT PLT'S 29/79


COR DEVEL. AXILLARY TUMOR (WAS AT. CARRIER)
PBT DEVEL. HEMANGIOMA (" ")

PHOSPHORUS BALANCE, STYLAB-2



URINARY ALDOSTERONE

CBB ○
 SPT ■
 FLT ▲




SKYLAB SUMMARY


PLASMA/SERUM BIOCHEMICAL CHANGES SUBSTANCE INCREASED

IN-FLIGHT	POST FLIGHT
POTASSIUM	POTASSIUM (R-D)
CALCIUM*	CALCIUM*
PHOSPHORUS*	PHOSPHORUS*
CREATININE	CREATININE
CORTISOL*	CORTISOL
ANGIOTENSIN I*	ANGIOTENSIN I
	THYROID STIMULATING HORMONE *
	THYRONINE *
	GLUCOSE*
	INSULIN
	ALDOSTERONE*
	CHOLESTEROL
	TRIGLYCERIDES
	S-GLUTAMIC OXALOACETIC
	TRANS
	CREATINE PHOSPHOKINASE
	TOTAL PROTEIN*
	CARBON DIOXIDE *

* P < 0.05



NITROGEN BALANCE, 1971-72



SKYLAB SUMMARY

PLASMA/SERUM BIOCHEMICAL CHANGES

SUBSTANCE/PROPERTY DECREASED	
IN-FLIGHT	POSTFLIGHT
SODIUM*	SODIUM
CHLORIDE*	CHLORIDE*
OSMOLALITY*	OSMOLALITY
GLUCOSE*	ADRENOCORTICOTROPIC HORMONE
INSULIN*	BLOOD UREA NITROGEN
ADRENOCORTICOTROPIC HORMONE*	CHOLESTEROL
ALDOSTERONE	URIC ACID
	MAGNESIUM
	LACTIC DEHYDROGENASE
	TOTAL BILIRUBIN
	ALBUMIN

* $P \leq 0.05$

SKYLAB CREW HEALTH TREND - BODY WEIGHT


PERCENT
CHANGE
FROM
PREFLIGHT
AVERAGE

DAYS OF MISSION


10 20 30 40 50 60 70 80

COMMANDER
SCIENTIST PILOT
PILOT







THE EFFECT OF PROLONGED BEDREST ON
MEAN URINARY CALCIUM EXCRETION
(3 SUBJECTS)




URINARY CORTISOL

DBB ○
 SP1 ■
 PLT ▲



AVERAGE LEG VOLUME CHANGE,
POSTFLIGHT - SKYLAB


URINARY CALCIUM, SKYLAB-4



HAAS-0-04-000

URINE VOLUME

CDR ○
SPT ■
PLT ▲



SKYLAB SUMMARY

24-HOUR URINE RESULTS


SUBSTANCE/PROPERTY INCREASED	
IN-FLIGHT	POSTFLIGHT
SODIUM	EPINEPHRINE
POTASSIUM	NOREPINEPHRINE
CHLORIDE	ALDOSTERONE
OSMOLALITY	CORTISOL
CALCIUM	TOTAL
PHOSPHORUS	17-KETOSTEROIDS
ALDOSTERONE	
CORTISOL	
17-KETOSTEROIDS	

RED CELL MASS

MEAN % CHANGE

SKYLAB	<u>2</u>	<u>3</u>	<u>4</u>
R + 0	-14.3	-12.2	-6.8
R + 14	-15.6	-7.5	-4.5
R + 42, 45	-5.2	+2.6	+3.7
R + 67	+0.2	-	-

CALCIUM BALANCE, STUDY-1




NR18-2-74-0888

URINARY ADH

COR. O

SPT ■

PLT ▲



SKYLAB SUMMARY

24-HOUR URINE RESULTS


SUBSTANCE/PROPERTY DECREASED	
IN-FLIGHT	POSTFLIGHT
ANTIDIURETIC HORMONE ±	SODIUM
EPINEPHRINE ±	POTASSIUM
NOREPINEPHRINE	CHLORIDE
TOTAL 17-HYDROXYCORTICOSTEROIDS ±	OSMOLALITY
URIC ACID	PHOSPHORUS
	MAGNESIUM
	URIC ACID
	ANTIDIURETIC HORMONE ±
	TOTAL 17-HYDROXYCORTICOSTEROIDS
	TOTAL 17-KETOSTEROIDS

MO92, LOWER BODY NEGATIVE PRESSURE CHANGE IN CALF VOLUME, SL-4 PLT




WASH-5-78-00001

MO92 LOWER BODY NEGATIVE PRESSURE SKYLAB 4




RED CELL MASS/SKYLAB

MISSION
DURATION

313 (a)


313 (a)

314 (a)



SAIR 3-74-4087

M092 LOWER BODY NEGATIVE PRESSURE SKYLAB 4



SKYLAB CREWS RETICULOCYTE COUNTS

PERCENTAGE OF PREMISSION MEAN

<u>DAY</u>	<u>MISSION</u>		
	<u>2</u>	<u>3</u>	<u>4</u>
R + 0	44	69	88
R + 1	53	-	117
R + 3	67	87	87
R + 7	80	184	143
R + 14	86	227	184
R + 21	93	234	180

HISTORY OF MOTIVATION AND VESTIBULOMETRIC FINDINGS IN THE NINE STRIKEAYS


STRIKEAY	AGE	HISTORY OF MOTIVATION								Ocular Function		HEALTH STATUS	GENERAL COMMENTS AND DATA
		SICKNESS		REASON FOR VISIT		TYPE OF VISIT		DATE OF VISIT	TIME OF VISIT	DATE OF VISIT	TIME OF VISIT		
		DATE	TIME	DATE	TIME	DATE	TIME						
1	52	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
2	53	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
3	54	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
4	55	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
5	56	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
6	57	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
7	58	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
8	59	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%
9	60	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	1/10/50	10:00	100%	100%

*ALL STRIKEAYS ON THIS DATE


*ALL STRIKEAYS ON THIS DATE

*ALL STRIKEAYS ON THIS DATE

M092 LOWER BODY NEGATIVE PRESSURE SKYLAB 4



THE LOWER BODY NEGATIVE PRESSURE PROTOCOL USED FOR SKYLAB CARDIOVASCULAR EVALUATIONS ASSESSING ORTHOSTATIC TOLERANCE




RED CELL MASS


<u>MISSIONS</u>	<u>MEAN % CHANGE</u>
GEMINI 5	-21
GEMINI 7	-14
APOLLO 7-8	- 2
APOLLO 14-17	-10 $\pm$ 1
SKYLAB 2	-14
SKYLAB 3	-12
SKYLAB 4	- 7
CONTROLS	-0.4


FREQUENCY DISTRIBUTION OF RETINA THICKNESS SUSCEPTIBILITY SCORES OF THE NINE SKYLAB ASTRONAUTS



MO92 LOWER BODY NEGATIVE PRESSURE SKYLAB 4








SCIENTIFIC CATEGORIES OF DIFFERENT LEVELS OF ACQUISITION STRESS

Category	Polysymptomatic 16 points	* Major 8 points	Minor 4 points	Residual 2 points	None ¹ 0 point
Respiratory symptoms (w/ or w/o)	Respiratory B/ wheezing	Respiratory B	Respiratory I	Episodic dyspnea	Episodic wheezing
Run		Factor II	Factor II	Factor I	Running/Polysymptomatic severity III
Cold sensitivity		III	II	I	
Increased salivation		III	II	I	
Prophylaxis		III	II	I	
Fast					Residual (percentage) III
Control recovery system					Control (percentage) Fast control III Fast speed III
Levels of Severity Identified by Total Points Scored					
From Extreme	Severe Malaise (0-10)	Moderate Malaise A (0-10)	Moderate Malaise B (0-10)	High Malaise (0-1)	
16 points	0 - 15 points	5 - 7 points	1 - 4 points	1 - 2 points	

*Agg. = seasonal (wheezing) symptoms

III = severe or marked II = Moderate, I = slight

M092 LOWER BODY NEGATIVE PRESSURE **SKYLAB 4**





MO92 LOWER BODY NEGATIVE PRESSURE

SKYLAB 4
PREFLIGHT

SP1


DOT 299






SKYLAB GENERAL SUMMATION

- BIOMEDICAL RESULTS SHOW THAT MAN CAN ADAPT AND FUNCTION EFFECTIVELY IN WEIGHTLESS ENVIRONMENT FOR EXTENDED PERIODS
- DAILY IN-FLIGHT PERSONAL EXERCISE REGIMENS COUPLED WITH APPROPRIATE DIETARY INTAKE AND PROGRAMED ADEQUATE SLEEP/WORK/RECREATION PERIODS ESSENTIAL FOR MAINTAINING CREW HEALTH AND WELL-BEING
- NO UNFAVORABLE PHYSIOLOGICAL CHANGES NOTED THAT WOULD PRECLUDE LONGER DURATION MANNED SPACE FLIGHTS; HOWEVER, RESEARCH REQUIRED TO UNDERSTAND THE MECHANISMS RESPONSIBLE FOR MANY OBSERVED CHANGES
- REMEDIAL OR PREVENTIVE MEASURES MAY BE REQUIRED FOR MISSION DURATIONS IN EXCESS OF 9 TO 12 MONTHS (E.G., BONE DEMINERALIZATION COUNTERMEASURES)
- IDEALLY, FURTHER OBSERVATIONS OF MAN IN EARTH ORBIT FOR AN UNINTERRUPTED PERIOD OF 6 MONTHS SHOULD PRECEDE A MARS-TYPE MISSION



NOTES: SUBJECTS: SYMPTOMATOLOGY OF 51.3 ASTRONAUTS



SECLAB 4 ANTIMOTION SICKNESS MEDICATION

MISSION EVENT	APPROX. TIME (HOURS C.S.T.)	CHWASER	SCIENTIST PILOT	PILOT
LAUNCH DAY (MO 1)				
AFTER INSERTION	0900		A	B
AFTER MO-2*	1100	A		
AFTER DOCKING	1700	B	B	B
	2300			B
MO-2 & MO-3	0800	A	A	B
ON ARRIVAL	1000	A	A	
	1400	B	B	B
MO-4	0800	A	A	B
ON ARRIVAL	1400			B
MO-6	RESTING		B	
MO-10	RESTING		B	
MO-82	RESTING		B	
MO-84		A	A	B
ABOUT 2 HOURS PRIOR TO SPLASH (ENTRY)				

A SCOPOLAMINE / DIMENHYDRATE 10.00/50.0 mg

B PROMETHAZINE / EPHEDRINE 125/50 mg


* FIRST PHASE MANEUVER

HEMATOLOGIC SYSTEM


- FINDING:
 - RED CELL MASS LOSS
- ETIOLOGICAL FACTOR(S)
 - INCREASED DESTRUCTION - NO EVIDENCE
 - MARROW SUPPRESSION
 - ? TOXIC
 - ? PHYSICAL (INCREASED BLOOD VOLUME, MARROW VENOUS PRESSURE)
- CANDIDATE FUTURE STUDIES
 - GROUND-BASED MARROW-SUPPRESSION FACTORS
 - VALIDATE SKYLAB RESULTS ON LONGER EARTH-ORBITAL FLIGHTS

LIFE SCIENCES CORE CONCEPT

(COMMON OPERATIONS RESEARCH EQUIPMENT)



MOTION SICKNESS SYMPTOMATOLOGY OF SL-4 ASTRONAUTS



SENSITIVITY TO ACTION DECISION UNDER OPERATIONAL CONDITIONS

EXPERIMENT	N	STANDARD TEST								STANDARD TEST	STANDARD TEST			STANDARD TEST
		STANDARD TEST		STANDARD TEST		STANDARD TEST		STANDARD TEST			STANDARD TEST			
		STANDARD TEST	STANDARD TEST	STANDARD TEST	STANDARD TEST	STANDARD TEST	STANDARD TEST	STANDARD TEST	STANDARD TEST		STANDARD TEST	STANDARD TEST		
1	1	1	1	1	1	1	1	1	1	1	1	1	1	
2	2	2	2	2	2	2	2	2	2	2	2	2	2	
3	3	3	3	3	3	3	3	3	3	3	3	3	3	
4	4	4	4	4	4	4	4	4	4	4	4	4	4	
5	5	5	5	5	5	5	5	5	5	5	5	5	5	
6	6	6	6	6	6	6	6	6	6	6	6	6	6	
7	7	7	7	7	7	7	7	7	7	7	7	7	7	
8	8	8	8	8	8	8	8	8	8	8	8	8	8	
9	9	9	9	9	9	9	9	9	9	9	9	9	9	
10	10	10	10	10	10	10	10	10	10	10	10	10	10	

1. EXPERIMENT 1: (ACTION DECISION) ACTION DECISION

2. ACTION DECISION

3. ACTION DECISION

4. ACTION DECISION

5. ACTION DECISION

6. ACTION DECISION

7. ACTION DECISION

8. ACTION DECISION

9. ACTION DECISION

10. ACTION DECISION

1. EXPERIMENT 1: (ACTION DECISION) ACTION DECISION

2. ACTION DECISION

3. ACTION DECISION

4. ACTION DECISION

5. ACTION DECISION

6. ACTION DECISION

7. ACTION DECISION

8. ACTION DECISION

9. ACTION DECISION

10. ACTION DECISION

SKYLAB 2 (28 DAY MISSION)


RED CELL MASS % CHANGE

	<u>CDR</u>	<u>SPT</u>	<u>PLT</u>
R + 0	-15.2	-15.6	-12.1
R + 13	-17.5	-16.4	-12.8
R + 42	- 8.2		- 2.2
R + 67	-3.0	+ 1.5	+ 2.0

CARDIOVASCULAR SYSTEM (CONCL)

- RULE OUT
 - PERMANENT MYOCARDIAL DAMAGE (CELLULAR LEVEL) - REMOTE
- CANDIDATE FUTURE CARDIOVASCULAR STUDIES
 - IN-DEPTH, NONINVASIVE CARDIOVASCULAR DYNAMICS MONITORING
 - INVASIVE PRESSURE/VOLUME/FLOW CHANGES IN EARLY FLIGHT (ANIMAL)
 - DEMONSTRATE PRESENCE OR ABSENCE OF GAUQUER-HENRY REFLEX
 - TOTAL BODY EXERCISE REGIMEN TO MAINTAIN INTEGRITY OF ANTI-GRAVITY AS WELL AS MAJOR MUSCLE GROUPS
 - ASSESS ROLE OF VENOUS (CAPACITANCE) VESSELS IN OBSERVED DECONDITIONING PROCESS
 - ASSESS ROLE OF FATIGUE

THE ROTATING LITTER CHAIR (RLC) IN THE POSTURE ASSESSMENT TEST WOOD



SKYLAB 3 (59 DAY MISSION)

RED CELL MASS % CHANGE

	<u>CDR</u>	<u>SPT</u>	<u>PLT</u>
R + 0	-6.1	-19.8	-10.6
R + 14	-2.7	-13.8	-5.9
R + 45	+3.0	+1.7	+3.1

CARDIOVASCULAR SYSTEM


◆ FINDINGS

- ◆ POSTFLIGHT ORTHOSTATIC INTOLERANCE
- ◆ POSTFLIGHT DIMINISHED EXERCISE CAPACITY

◆ PROBABLE ETIOLOGICAL FACTORS

- ◆ DECREASED EFFECTIVE CIRCULATING BLOOD VOLUME
POSTFLIGHT
- ◆ DIMINISHED VENOUS RETURN AT ONE g
- ◆ MUSCULAR IMBALANCE CAUSED BY FUNCTIONAL DISUSE
ATROPHY OF ANTIGRAVITY MUSCLES
- ◆ ALTERED INTERNAL MILIEU (FLUID/ELECTROLYTE DYNAMIC
FLUX) DURING EARLY POSTFLIGHT PERIOD
- ◆ ? ALTERED VENOUS REFLEXES/TONE
- ◆ FATIGUE

MO92, LOWER BODY NEGATIVE PRESSURE CHANGE IN RESTING-CALF CIRCUMFERENCE, SL-4 CREW




SKYLAB 4 (84 DAY MISSION)

RED CELL MASS % CHANGE


	<u>CDR</u>	<u>SPT</u>	<u>PLT</u>
R + 0	-4.8	-8.3	-7.4
R + 14	-4.0	-3.9	-5.5
R + 31	+4.7	+2.3	+4.0

Motion Sickness Symptomatology in SL 1 of Astronauts




NASA-D-74-4837

DAMAGED ORBITAL WORKSHOP



ANTHROPOMETRIC CHANGES SL-4 PLT



SKYLAB

IRON KINETICS AT RECOVERY

	<u>CREW</u>	<u>CONTROL</u>
SERUM Fe	92	109
T 1/2	127	164
TURNOVER	0.32	0.30

SPECIAL IN-FLIGHT TESTS (ORIGINALLY APPROVED)

- ENVIRONMENTAL MICROBIOLOGY
- RADIATION MEASUREMENTS
- WATER POTABILITY
- CO₂/DEWPOINT MONITOR
- CREW MICROBIOLOGY
- ORAL HEALTH

SKYLAB CREWS AT RECOVERY

- NORMAL
 - RBC SURVIVAL
 - IRON KINETICS
- DECREASED
 - RED CELL MASS
 - RETICULOCYTES

¹⁴C-GLYCINE MEAN
RED CELL LIFE TIME
DAYS

MEAN ± S.E.

CREW 123 ± 2

CONTROL 121 ± 4

SKYLAB RBCM CHANGE THEORY

- INCREASED SPLENIC RETICULOENDOTHELIAL
ACTIVITY
- DECREASED PLASMA VOLUME
- INCREASED PHOSPHORUS
- NO ERYTHROPOIETIN CHANGE
- RELATIVE BONE MARROW INHIBITION

SKYLAB INFILIGHT MEDICAL SUPPORT SYSTEM - MMS SUBSYSTEMS

RESPIRATOR
THERM. EXCHANGER

RESPIRATORY
CONTAINER

MMS RITS

DRUGS SURVIV
PT. FLUIDS
THERMAL DRUGS
BOTTLES DRUGS
MICROBIOLOGIC
MINOR SURGERY KIT
MICROBIOLOGIC

HEMATOCENTRIF
HEMATOCENTRIF
DIAGNOSTIC
BANDAGE
BOTTLES
THERMOCENTRIF




MEAN ^{51}Cr RED CELL HALFTIMES

DAYS

	<u>CREW</u>	<u>CONTROL</u>
PRE	26.1	24.2
POST	24.2	22.9
DELTA	-1.9	-1.3

SKYLAB BLOOD SAMPLE PROCESSOR



BLOOD DRAWN SKYLAB MISSIONS

<u>MISSION/DAYS</u>		<u>TOTAL ml</u>	<u>MEAN ml/DAY</u>
SL2	28	794	10
SL3	59	805	8
SL4	84	889	6

SIGNIFICANT BIOMEDICAL FINDINGS IN GEMINI PROGRAM

- MODERATE LOSS OF RED CELL MASS
- MODERATE ORTHOSTATIC INTOLERANCE
- MODERATE LOSS OF EXERCISE CAPACITY
- MINIMAL LOSS OF BONE DENSITY
- MINIMAL LOSS OF CALCIUM AND MUSCLE
NITROGEN
- HIGH METABOLIC COST OF EXTRAVEHICULAR
ACTIVITY


FLUID/ELECTROLYTE AREA

- PRESKYLAB WORKING HYPOTHESES
- PARADOXICAL SKYLAB URINARY FINDINGS
 - ↓ ADH SECRETION (↓ SKYLAB 2)
 - ↓ ALDOSTERONE SECRETION
 - ↓ Na^+ K^+ EXCRETION
 - ↓ OSMOLALITY
 - ↓ CORTISOL
- INTERPRETATION
 - INTERNAL HOMOSTASES MAINTAINED
 - RENAL FUNCTION MAINTAINED
 - INTERACTION OF COMPLEX HUMORAL (AND THERMODYNAMIC) FACTORS REQUIRED TO MAINTAIN HOMOSTASES DURING WEIGHTLESSNESS
- FUTURE STUDIES:
 - DEMONSTRATE GAUER-HENRY REFLEX
 - RENAL HEMODYNAMICS IN ZERO g
 - RENAL RESPONSE TO WATER/SALT LOADS, DEHYDRATION IN ZERO g
 - HUMORAL INTERACTIONS INVOLVED IN ABOVE

SKYLAB MINERAL/CALORIC, FLUID/ELECTROLYTE SUMMARY

- MODERATE LOSSES OF CALCIUM, PHOSPHORUS, AND NITROGEN HAVE BEEN OBSERVED COMPARABLE TO THOSE SEEN IN BED-RESTED SUBJECTS: 6 GRAMS/MONTH CALCIUM OR 0.6 PERCENT/MONTH TOTAL BODY CALCIUM
- RATE OF CALCIUM LOSS WOULD NOT PRECLUDE EXTENDED MISSIONS OF 6 TO 9 MONTHS' DURATION. LONGER MISSIONS MAY REQUIRE REMEDIAL MEASURES
- SIGNIFICANT 0.6 CALCIUM MINERAL LOSS:
SKYLAB 3: SPT: 7.4 PERCENT
SKYLAB 4: SPT: 4.5 PERCENT; PLT: 7.8 PERCENT
- CALORIC REQUIREMENTS DURING FLIGHT IDENTICAL WITH INDIVIDUAL REQUIREMENTS AT ONE G
- RENAL FUNCTION UNIMPAIRED; HOWEVER, APPARENTLY UNIQUE ADAPTIVE FUNCTIONAL CHANGES OBSERVED REQUIRE FURTHER STUDY
- SKYLAB 4 ANTHROPOMETRIC STUDIES CONSISTENT WITH PREDICTED SHIFT OF BODY FLUIDS DURING WEIGHTLESS FLIGHT

CARDIAC OUTPUT DURING SUBMAXIMAL EXERCISE




MUSCULOSKELETAL SYSTEM

- FINDING
 - MODERATE LOSSES OF CALCIUM, NITROGEN, AND PHOSPHORUS
- POSSIBLE ETIOLOGICAL FACTORS
 - PRIMARY - LOSS OF GRAVITY GRADIENT
 - SECONDARY
 - ABSENCE OF WEIGHT BEARING
 - ABSENCE OF HYDROSTATIC VENOUS GRADIENT
 - ? HORMONAL IMBALANCE (ELEVATED CORTISOL SECRETION)
 - COMBINATIONS OF ABOVE
- CANDIDATE FUTURE STUDIES
 - ABSOLUTE CATABOLIC IN-FLIGHT CHANGES (BONE, MUSCLES) IN ANIMALS
 - COUNTERMEASURE EVALUATION
 - DIETARY
 - PHYSICAL
 - ? HORMONAL
 - SELECTION CRITERIA FOR PROLONGED MISSIONS (PREDICTION FORMULA)

STROKE VOLUME DURING SUBMAXIMAL EXERCISE





SKYLAB NEUROPHYSIOLOGICAL/ PERFORMANCE SUMMARY

- AFTER INITIAL IN-FLIGHT ADAPTIVE PERIOD, ALL CREWMEN SHOW MARKED INCREASED TOLERANCE TO MOTION SICKNESS MEASURED BY ROTATION AND HEAD MOVEMENTS. TOLERANCE THRESHOLD GRADUALLY REVERTS TO BASELINE POSTFLIGHT. ANTI-MOTION MEDICATIONS HELPFUL IN CONTROL OF SYMPTOMS.
- NO MAJOR DISTURBANCES NOTED IN QUANTITY OR QUALITY OF SLEEP.
- CAREFULLY REGULATED CREW WORK/REST CYCLES ESSENTIAL FOR MAINTAINING CREW EFFICIENCY.
- POSTFLIGHT HYPERREFLEXIA CONFIRMED AND QUANTITATED.
- IMPROVED IN-FLIGHT PERFORMANCE EFFICIENCY EXHIBITED BY ALL CREWS.
- OVERLEARNING RECOMMENDED FOR CRITICAL TASKS ON SHORT-DURATION MISSIONS.

SKYLAB HEMATOLOGY/IMMUNOLOGY/ CYTOLOGY SUMMARY


- LOSS OF RED CELL MASS OBSERVED POSTFLIGHT - APPEARS TO BE SUPPRESSION OF RED CELL PRODUCTION RATHER THAN DESTRUCTION
- RED BLOOD CELL MASS LOSS NOT RELATED TO MISSION DURATION
- NO SIGNIFICANT CHANGES OBSERVED IN HUMORAL OR CELLULAR IMMUNE RESPONSES
- CELLULAR PROLIFERATION (TISSUE CULTURE) NORMAL DURING SPACE FLIGHT

DAILY PERSONAL EXERCISE PROTOCOL SELECTED BY EACH SL-4 CREWMAN

EXERCISE	CDR	SPT	PLT
LEG ERGOMETRY (WATT-MIN)	5000	8337	6000
MINISYM (TOTAL REPETITIONS)	100	200	200
SPRINGS (TOTAL REPETITIONS)	75	0	120
TORSO ISOMETRICS (TOTAL REPETITIONS)	30	0	20
TREADMILL			
WALK (MIN)	10	0	10
RUN (MIN)	1	0	0
SPRINGS (REPETITIONS)	300	1000	100
TOE RISES (REPETITIONS)	200	200	75

ADAPTIVE RESPONSES TO WEIGHTLESSNESS-PRE Skylab Hypothesis





SKYLAB VESTIBULAR FINDINGS

- POSSIBLE ETIOLOGICAL FACTORS

- OTOLITH RECEPTOR PHYSIOLOGICAL DEAFFERENTATION

- † MODULATING INFLUENCE ON CANALS

- DIRECT INTERACTION WITH CANALS

- RULE OUT:

- INFLUENCE OF HYPERVOLEMIA


- (TRANSIENT † VENOUS PRESSURE) ON VESTIBULAR SYSTEM

NASA-SP-75-4881

ORBITAL WORKSHOP WITH THERMAL SHIELDS DEPLOYED




METABOLIC ANALYZER FUNCTIONAL SCHEMATIC



NASA-S-81-208-X

SKYLAB



SKYLAB VESTIBULAR FINDINGS (CONCL)

• CANDIDATE FUTURE STUDIES

- ROLE OF ALTERED CUES: VISUAL, KINESTHETIC OTHER SENSORY
- EFFECT OF OVERHYDRATION, DEHYDRATION AND INCREASED VENOUS PRESSURE ON MOTION SICKNESS THRESHOLD
- PREDICTIVE TEST FOR ZERO g SPACE SICKNESS SUSCEPTIBILITY (? PARABOLIC FLIGHTS)
- BASIC STUDIES REGARDING ETIOLOGY
- ROLE OF ONE g TRAINING IN PREVENTION
- IMPROVED MEDICATIONS FOR PREVENTION/CONTROL

SKYLAB CARDIOVASCULAR SUMMARY

- CARDIOVASCULAR DISORIENTATION WAS OBSERVED DURING FLIGHT; CHANGES APPEAR ADAPTIVE IN NATURE AND TEND TO STABILIZE AFTER 4 TO 6 WEEKS
- CARDIOVASCULAR CHANGES DO NOT IMPAIR CREW HEALTH OR ABILITY TO FUNCTION EFFECTIVELY IN WEIGHTLESS FLIGHT
- LOWER-BODY RELATIVE-PRESSURE TESTS PROVIDE FAIRLY RELIABLE PREDICTIVE INDEX OF POSTFLIGHT CARDIOVASCULAR STATUS
- CARDIAC ELECTRICAL ACTIVITY AS MEASURED BY VECTOR-CARDOGRAM WAS NOT SIGNIFICANTLY ALTERED AND REMAINED WITHIN PHYSIOLOGICAL LIMITS
- DECREASED CARDIAC OUTPUT NOTED IN CREWMEN POSTFLIGHT; THOUGHT TO BE RELATED TO REDUCED VENOUS RETURN
- SINGLE EPISODE OF SIGNIFICANT CARDIAC ARRHYTHMIA IN ONE SKYLAB 3 CREWMAN DURING EXERCISE EARLY IN MISSION
- NO SIGNIFICANT IN-FLIGHT DECREMENT IN WORK CAPACITY OR PHYSIOLOGICAL RESPONSES TO EXERCISE
- ALL CREWMEN HAVE SHOWN POSTFLIGHT INCREASE IN EXERCISE CAPACITY AND ALTERED PHYSIOLOGICAL RESPONSES
- SKYLAB 3 AND 4 CREWS RETURNED TO PREFLIGHT CARDIOVASCULAR STATUS BY 7TH AND 14TH SKYLAB 3 CREW AT 7TH; ENHANCED EXERCISE BY SKYLAB 3 AND 4 CREWMEN THOUGHT TO BE FACTOR IN IMPROVED RECOVERY RATE

M171 EXPERIMENT PROTOCOL

5 MINUTES

REST

5 MINUTES

25 PERCENT OF MAXIMUM $\dot{V}O_2$

5 MINUTES

50 PERCENT OF MAXIMUM $\dot{V}O_2$

5 MINUTES

75 PERCENT OF MAXIMUM $\dot{V}O_2$

5 MINUTES

RECOVERY

3 CREWMEN

5 TIMES IN 28 DAY MISSION

8 TIMES IN 56 DAY MISSION