## The MANDRILL\*, or Ribbed-Nofe BABOON.

THE uglines of this baboon is perfectly disjoining. His node, or rather his two nothin, are flat, from which a fnot perpetually runs, and he licks it into his mouth with his tongue. His head is very large, and his muzzle long. His body is fupua, and his buttecks are of a blood colour. His amus is confipienous, and fituated almost as high as the loins. His face is of a violet colour, and furrounded on each fide with deep longitudinal wrinkles, which

• Ribbed-nofe baboon, with a long raiked nofe compreffed fidewife, of a purple colour, and ribbed obliquely on each file; on the chie, a fantr picked orange beard; sail very hairy, about two inches long, which it carries erect; buttocks naked; hair foft, daily mottled with yellow; longth, from node to tail, about two fett, Pennaer's "somet' of Space", by 1-101.

Simia maiuson, candata, subbarbata, genis coroleis striatis, natibus calvis; Liner. Sys. Nat. p. 35.

Cercopithecus cynocephalus, parte corporis anteriore longis

pills obfits, nato violacco nudo; Briffen. Regn. Anim. p. 214.

Maudrill, the name which the British, who frequent the coaft
and Guiner, have given to this animal.

There is a firange fort of animal, called by the white men in Guiney, a manuful! to the white it is to called, I know not, nor did I ever hear of the name before i neither can though one who call them fo tell, except it be for its near refemblance of a human creature, though nothing at all like an ape; Snith's Feynge to Guiney, 6, 62.

VOL. VIII. 1 augment

augment the fullenness and deformity of his afpect. He is likewise larger, and perhaps ffronger, than the absoors but at the fame time, he is more peaceable; and less ferocious. We here give figures of both the male and the formals, which we have feen alive. Whether they had received a better education, or if they be nutrally more gentle than the absoon, they appeared to be more trackable and less impudent; but they were equally differeable.

This species of baboon is found on the Gold Coast, and in the other fouthern provinces of Africa, where he is called begge by the Negroe, and mandrill by the Europeans. Next to the orang-outang, he is the largest of all the aps or baboons. Smith relates \*, that he had a pre-

\*\* The body of the manifold, when full grown, is a kipse circumstrear as a sulfaction stars. It his grow earth sheen, and his feet langer, his name and handle in proportion. The shad the free independent of the first star and handle in proportion. The shad handle is supported to the star of the shad handle in the first him. The face, which is covered with a shad handle in the shad handle in

fent of a female mandrill, which was only fix months old, and that it was as large as an adult baboon. He adds, that these mandrills walk always on two feet; that they weep and groan like men; that they have a violent passion for women, which they never fail to gratify when they find a woman at a distance from relief.

Distinctive Characters of this Species.

The mandrill has check-pouches, and callofities on the butches. The tail exceeds not two or three inches. The canine teeth are much thicker and longer than thole of man. The muzzle is very thick, very long, and furrowed on each fide with deep longitudinal wrinkles. The face is naked, and of a blueifs closur. The cars, as well as the palm of the hands and foles of the feet, are naked. The hair is long.

Smill's Piggger to Galage, p. 5, 1. In the fine country, this saimal is called deep or degree, and markell', and they likewise client the cauge, country, page and deall. There is much failinity in their market, and they are probably deviated from each other; And, in fall, the pages and logge, or the drill and mandrill, have feveral common charders. But he fall is an age without a mill, attend nabed, and with a fast over farce; and the forced is a balance with attitle and English Imagings, guides san as ig a personal and deall, in the jurgen of fonce of our French provinces, decested a sejement likewise. of a reddish brown upon the body, and gray upon the breast and belly. He walks on two feet offener than on four—When erect, he is four or four and a half feet high; and some of them seem to be still larger. The semales are subject to the menser.



FEMALE MANDRILL.